

Metric Lie Algebras with Maximal Isotropic Centre

Ines Kath and Martin Olbrich

February 1, 2008

Abstract

We investigate a certain class of solvable metric Lie algebras. For this purpose a theory of twofold extensions associated to an orthogonal representation of an abelian Lie algebra is developed. Among other things, we obtain a classification scheme for indecomposable metric Lie algebras with maximal isotropic centre and the classification of metric Lie algebras of index 2.

Contents

1	Introduction	1
2	A structure theorem	5
3	Equivalence of twofold extensions and the isomorphism problem	10
4	A decomposability criterion	17
5	The case of maximal isotropic centre	20
6	Metric Lie algebras of index 2	27

1 Introduction

In the present paper we study a certain class of metric Lie algebras in a systematic way. Here a metric Lie algebra is a finite-dimensional real Lie algebra equipped with an invariant non-degenerate symmetric bilinear form. An isomorphism of metric Lie algebras is by definition a Lie algebra isomorphism which is in addition an isometry with respect to the given inner products.

A metric Lie algebra $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is called decomposable if it contains a proper ideal $\mathfrak{i}$ which is non-degenerate (i.e. $\langle \cdot, \cdot \rangle|_{\mathfrak{i} \times \mathfrak{i}}$ is non-degenerate), and indecomposable otherwise. Then any metric Lie algebra is the orthogonal direct sum of indecomposable metric Lie algebras. Thus, in order to understand metric Lie algebras one has to understand the indecomposable ones, and the natural (but certainly too ambitious) task would be to obtain a classification of indecomposable metric Lie algebras up to isomorphism.

Of course, in the case of definite inner products there is nothing left to do. Any indecomposable metric Lie algebra with a definite inner product is either compact simple or one-dimensional. However, the situation changes dramatically if we consider indefinite inner products. Here the simple and one-dimensional Lie algebras constitute only a small part of the set of all indecomposable metric Lie algebras.

In [MR 85] A. Medina and Ph. Revey proved a first structure result. They considered the following construction, which starts with a metric Lie algebra and produces metric Lie algebras of higher dimension.

Let $(\mathfrak{b}, \langle \cdot, \cdot \rangle_{\mathfrak{b}})$ be a metric Lie algebra, $\mathfrak{l}$ a Lie algebra with invariant (possibly degenerate) symmetric bilinear form $\langle \cdot, \cdot \rangle_{\mathfrak{l}}$ and $\rho : \mathfrak{l} \rightarrow \text{Der}_{\mathfrak{a}}(\mathfrak{b}, \langle \cdot, \cdot \rangle_{\mathfrak{b}})$ a representation of $\mathfrak{l}$ by antisymmetric derivations on $(\mathfrak{b}, \langle \cdot, \cdot \rangle_{\mathfrak{b}})$. Let $\mathfrak{d}_{\rho}(\mathfrak{b}, \mathfrak{l})$ be the extension

$$0 \longrightarrow \mathfrak{l}^* \longrightarrow \mathfrak{d}_{\rho}(\mathfrak{b}, \mathfrak{l}) \longrightarrow \mathfrak{b} \rtimes_{\rho} \mathfrak{l} \longrightarrow 0 \quad (1)$$

of the semi-direct sum of $\mathfrak{b}$ and $\mathfrak{l}$ defined by the coadjoint representation of $\mathfrak{l}^*$ on the second factor of $\mathfrak{b} \rtimes_{\rho} \mathfrak{l}$ and the cocycle $\beta \in Z^2(\mathfrak{b} \rtimes_{\rho} \mathfrak{l}, \mathfrak{l}^*)$ which is given by $\beta(X_1 + L_1, X_2 + L_2) = \langle X_1, \rho(\cdot)X_2 \rangle_{\mathfrak{b}}$ for $X_i \in \mathfrak{b}$ and $L_i \in \mathfrak{l}$, $i = 1, 2$. Define an inner product $\langle \cdot, \cdot \rangle$ on $\mathfrak{d}_{\rho}(\mathfrak{b}, \mathfrak{l})$ by

$$\langle \alpha_1 + X_1 + L_1, \alpha_2 + X_2 + L_2 \rangle = \langle X_1, X_2 \rangle_{\mathfrak{b}} + \langle L_1, L_2 \rangle_{\mathfrak{l}} + \alpha_1(L_2) + \alpha_2(L_1)$$

for $\alpha_i \in \mathfrak{l}^*$, $X_i \in \mathfrak{b}$ and $L_i \in \mathfrak{l}$, $i = 1, 2$. Then $(\mathfrak{d}, \langle \cdot, \cdot \rangle)$ is a metric Lie algebra and it is called a double extension of $\mathfrak{b}$ by $\mathfrak{l}$.

What Medina and Revey found in [MR 85] is that any non-simple non-abelian indecomposable metric Lie algebra $(\mathfrak{d}, \langle \cdot, \cdot \rangle)$ is isomorphic to a double extension $\mathfrak{d}_{\rho}(\mathfrak{b}, \mathfrak{l})$, where $\mathfrak{l}$ is one-dimensional or simple. The idea of the proof is to choose a minimal abelian ideal $\mathfrak{i} \neq 0$ in $\mathfrak{d}$ and to take $\mathfrak{l} = \mathfrak{d}/\mathfrak{i}^{\perp}$, $\mathfrak{b} = \mathfrak{i}^{\perp}/\mathfrak{i}$, $\mathfrak{l}^* = \mathfrak{i}$. See also [FS 96] for a nice presentation of the result and its applications to conformal field theory.

This structure result is very suitable for the classification of non-simple indecomposable Lorentzian Lie algebras (i.e. metric Lie algebras of signature $(1, q)$). The centre $\mathfrak{z}$ of such a Lie algebra is non-trivial. Furthermore, it is isotropic because of indecomposability. Hence, in this case $\mathfrak{i} = \mathfrak{z}$ is a canonical choice for a (one-dimensional) minimal abelian ideal. In particular, the Lie algebra is canonically isomorphic to a double extension. This makes it easy to study decomposability and to solve the isomorphism problem. The classification result is due to A. Medina ([M 85], see also Section 5 for an exact formulation of the result). Originally this result was not a consequence of the structure theorem, but the structure theorem arose as a generalization of its proof.

In [MR 85] Medina and Revey applied the structure theorem also for non-simple indecomposable metric Lie algebras of signature $(2, q)$. This yields a description of such Lie algebras as double extensions of (the already classified) Lorentzian Lie algebras by one-dimensional ones. Furthermore, Medina and Revey determined the (outer) antisymmetric derivations of a Lorentzian Lie algebra. So we know in principle how a non-simple indecomposable metric Lie algebras of signature $(2, q)$ looks like. However, this is not a classification. On one hand a metric Lie algebra of signature $(2, q)$ which is the double extension of a Lorentzian Lie algebra by a one-dimensional one can be decomposable and Medina and Revey did not study which of the obtained Lie algebras are really indecomposable. On the other hand it is not clear which of the obtained Lie

algebras are isomorphic. Contrary to the Lorentzian case the description of a metric Lie algebra of signature $(2, q)$ as a double extension is not canonical. So one can obtain isomorphic Lie algebras of signature $(2, q)$ even if one starts with non-isomorphic Lorentzian Lie algebras. Proposition 2.2 of the present paper is more adapted to this situation. It contains only canonical choices and allows a better study of indecomposability and isomorphy. In particular, it yields (together with Proposition 2.4 and the results of Section 5) a complete classification of indecomposable metric Lie algebras of signature $(2, q)$.

The classification result for signature $(2, q)$ was already announced in [BK 02] and it was one of our motivations for writing this paper to present a proof of this result in a quite systematic framework. Besides the classification for signature $(2, q)$ one can also find a classification of indecomposable metric Lie algebras of dimension $n \leq 6$ in [BK 02] although the paper does not concentrate on classification results. The emphasis is more on the construction of Lie groups with pseudo-Riemannian bi-invariant metric which admits parallel spinor fields using double extensions. One series of examples (called multiple extensions) corresponds to metric Lie algebras having a maximal isotropic centre and was constructed inductively by double extensions by a one-dimensional Lie algebra starting with an abelian Euclidean Lie algebra. The treatment of these examples suggested that there should be a better description of metric Lie algebras with maximal isotropic centre than the description by double extensions. This was a further motivation for writing the present paper.

From a geometric point of view, one is interested in the underlying semi-Riemannian symmetric space defined by a Lie group with bi-invariant metric. Of course, Lie groups are very special symmetric spaces, and one would like to understand all of them. On the other hand, any symmetric space is a homogeneous space of a distinguished Lie group, its transvection group. Let us discuss the relation between metric Lie algebras and semi-Riemannian symmetric spaces in little more detail.

Up to local isometry, a semi-Riemannian symmetric space is given by a triple $(\mathfrak{g}, \langle \cdot, \cdot \rangle, \sigma)$, where $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is a metric Lie algebra and σ is an involutive automorphism of $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ such that $\mathfrak{g} = [\mathfrak{m}, \mathfrak{m}] \oplus \mathfrak{m}$, where $\mathfrak{m}$ is the -1 -eigenspace of σ . By abuse of notation let us call such a triple simply a symmetric space. The signature of a symmetric space is by definition the signature of the restriction of $\langle \cdot, \cdot \rangle$ to $\mathfrak{m}$.

This description already supports the philosophy, which might be considered as the main motivation for a geometer to deal with metric Lie algebras, that any good method of investigation of metric Lie algebras should have a refinement which brings the automorphism σ into play, and then yields corresponding results for symmetric spaces. Thus, the first step in the study of symmetric spaces should be the study of metric Lie algebras.

Though symmetric spaces can be considered as more complicated objects than metric Lie algebras, in many respects one knows more about symmetric spaces than about metric Lie algebras. For instance, there is no thorough treatment of metric Lie algebras comparable to [CP 80]. Moreover, the classification of indecomposable symmetric spaces of index at most 2 is known for more than 30 years [CW 70], [CP 70] (see [N 02] for some necessary corrections). While the classification result for Lorentzian Lie algebras which was discussed above is already considerably younger, the corresponding result for metric Lie algebras of index 2 has not appeared in the literature before the

announcement given in [BK 02].

These circumstances are not only an accident. The study of metric Lie algebras cannot be reduced to that of symmetric spaces. The underlying symmetric space of (a Lie group associated to) a metric Lie algebra $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is the triple $(\tilde{\mathfrak{g}}, \langle \cdot, \cdot \rangle_{\tilde{\mathfrak{g}}}, \sigma)$, where

$$\tilde{\mathfrak{g}} = \{[(X, Y)] \in (\mathfrak{g} \oplus \mathfrak{g}) / \text{diag}(\mathfrak{z} \cap \mathfrak{g}') \mid X + Y \in \mathfrak{g}'\},$$

$\sigma([(X, Y)]) = [(Y, X)]$, and $\langle \cdot, \cdot \rangle_{\tilde{\mathfrak{g}}}$ is induced by the form $\langle \cdot, \cdot \rangle \oplus \langle \cdot, \cdot \rangle$ on $\mathfrak{g} \oplus \mathfrak{g}$. Here $\mathfrak{z}$ denotes the centre of $\mathfrak{g}$. We see that it is impossible to recover $\mathfrak{g}$ from its underlying symmetric space in general. What can be recovered is the scalar product $\langle \cdot, \cdot \rangle$ and the Lie algebra structure on $\mathfrak{g}/(\mathfrak{z} \cap \mathfrak{g}')$, only. Thus non-isomorphic metric Lie algebras may correspond to isomorphic symmetric spaces. Moreover, an indecomposable metric Lie algebra may give rise to a decomposable symmetric space.

Quite recently we learned that Berard Bergery and his collaborators in Nancy study symmetric spaces (in up to now unpublished work) based on a construction which we call twofold extension. This construction can be applied to metric Lie algebras as well and then becomes a close relative of the double extension described above. Roughly speaking, one simply replaces the semi-direct sum $\mathfrak{d}_\rho(\mathfrak{b}, \mathfrak{l})$ in (1) by a quite arbitrary extension of $\mathfrak{l}$ by $\mathfrak{b}$. In addition, we require $\mathfrak{b}$ to be abelian (therefore denoted by $\mathfrak{a}$ in the following). For this one has to pay the price that the Lie algebra $\mathfrak{l}$ can be more general than one-dimensional or simple. We feel that this method has the potential to give much more insight into the structure of general metric Lie algebras than the double extension method. In particular, it can be shown that any indecomposable non-simple metric Lie algebra can be written as such a twofold extension in a canonical way.

In the present paper we test this method in the simplest case, namely for abelian $\mathfrak{l}$. The precise construction of twofold extensions in this case is given in Proposition 2.1. It turns out that a relatively broad class of metric Lie algebras is already covered by this simple construction, namely those metric Lie algebras satisfying $\mathfrak{g}'' \subset \mathfrak{z}$ (Proposition 2.2). This class contains indecomposable metric Lie algebras with non-trivial maximal isotropic centre (Theorem 2.1) and indecomposable non-simple metric Lie algebras of index at most 2 (Proposition 2.4).

In Sections 3 and 4 we solve the crucial isomorphism and indecomposability problems for our twofold extensions. The key ingredient is the description of equivalence classes of twofold extensions associated to a fixed orthogonal representation $\rho : \mathfrak{l} \rightarrow \mathfrak{so}(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ by a certain cohomology set $\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a})$ (Corollary 3.1). This result is completely analogous to the classical identification of equivalence classes of extensions of a Lie algebra $\mathfrak{l}$ by an $\mathfrak{l}$ -module $\mathfrak{a}$ with the Lie algebra cohomology group $H^2(\mathfrak{l}, \mathfrak{a})$. Eventually, we can identify the set of isomorphism classes of indecomposable metric Lie algebras in our class with the union of orbit spaces of the actions of certain linear groups on sets made of the “indecomposable points” of the cohomology sets $\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a})$ (Proposition 4.2).

How the classification scheme provided by Proposition 4.2 can be made much more explicit in the case of maximal isotropic centre is discussed in Section 5. In particular, we give a complete classification of indecomposable metric Lie algebras having an at most 3-dimensional maximal isotropic centre in Theorem 5.1. Eventually, the classification of indecomposable metric Lie algebras of index 2 is presented in Theorem 6.1.

After we had obtained the results of Section 3 we realized that a nonlinear cohomology theory for orthogonal representations of general Lie algebras has been already developed by Grishkov in [Gr 98]. His results suggest that at least the results of Section 3 should have a direct analogue for non-abelian $\mathfrak{l}$. This and related topics might be the content of a forthcoming paper.

2 A structure theorem

We start this section by giving a construction method of certain metric Lie algebras.

Proposition 2.1 *Let $(\rho, \mathfrak{a})$ be an orthogonal representation of an abelian Lie algebra $\mathfrak{l}$ on the semi-Euclidean vector space $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$. Furthermore, choose a 3-form $\gamma \in \wedge^3 \mathfrak{l}^*$ and a cocycle $\alpha \in Z^2(\mathfrak{l}, \mathfrak{a})$ satisfying*

$$\langle \alpha(L_1, L_2), \alpha(L_3, L_4) \rangle + \langle \alpha(L_2, L_3), \alpha(L_1, L_4) \rangle + \langle \alpha(L_3, L_1), \alpha(L_2, L_4) \rangle = 0 \quad (2)$$

for all $L_1, L_2, L_3, L_4 \in \mathfrak{l}$. Then the bilinear map

$$[\cdot, \cdot] : (\mathfrak{l}^* \oplus \mathfrak{a} \oplus \mathfrak{l})^2 \longrightarrow \mathfrak{l}^* \oplus \mathfrak{a} \oplus \mathfrak{l}$$

defined by

$$\begin{aligned} \mathfrak{l}^* &\subset \mathfrak{z}(\mathfrak{l}^* \oplus \mathfrak{a} \oplus \mathfrak{l}) \\ [A_1, A_2] &= \langle \rho(\cdot) A_1, A_2 \rangle \in \mathfrak{l}^* \\ [A, L] &= \langle A, \alpha(L, \cdot) \rangle - L(A) \in \mathfrak{l}^* + \mathfrak{a} \\ [L_1, L_2] &= \gamma(L_1, L_2, \cdot) + \alpha(L_1, L_2) \in \mathfrak{l}^* + \mathfrak{a} \end{aligned}$$

for all $L, L_1, L_2 \in \mathfrak{l}$, $A, A_1, A_2 \in \mathfrak{a}$ is a Lie bracket on $\mathfrak{l}^* \oplus \mathfrak{a} \oplus \mathfrak{l}$ and the bilinear form $\langle \cdot, \cdot \rangle$ on $\mathfrak{l}^* \oplus \mathfrak{a} \oplus \mathfrak{l}$ defined by

$$\langle Z_1 + A_1 + L_1, Z_2 + A_2 + L_2 \rangle = \langle A_1, A_2 \rangle_{\mathfrak{a}} + Z_1(L_2) + Z_2(L_1)$$

for all $Z_1, Z_2 \in \mathfrak{l}^*$, $A_1, A_2 \in \mathfrak{a}$ and $L_1, L_2 \in \mathfrak{l}$ is an ad-invariant (non-degenerate) inner product on $\mathfrak{l}^* \oplus \mathfrak{a} \oplus \mathfrak{l}$.

Definition 2.1 *We denote the metric Lie algebra $(\mathfrak{l}^* \oplus \mathfrak{a} \oplus \mathfrak{l}, \langle \cdot, \cdot \rangle)$ constructed above by $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ and call it a twofold extension.*

Already in the case $\mathfrak{a} = 0$ this construction leads to interesting metric Lie algebras. For formal reasons we also include the trivial case $\mathfrak{l} = 0$ into our considerations.

Proof of Proposition 2.1. Consider the extension

$$0 \longrightarrow \mathfrak{a} \longrightarrow \mathfrak{d}_0 \longrightarrow \mathfrak{l} \longrightarrow 0$$

defined by ρ and $\alpha \in Z^2(\mathfrak{l}, \mathfrak{a})$. Now define a 2-form β on $\mathfrak{d}_0 = \mathfrak{a} \oplus \mathfrak{l}$ with values in $\mathfrak{l}^*$ by

$$\begin{aligned} \beta(A, L_1)(L_2) &= \langle A, \alpha(L_1, L_2) \rangle_{\mathfrak{a}} \\ \beta(A_1, A_2)(L) &= \langle L(A_1), A_2 \rangle_{\mathfrak{a}} \\ \beta(L_1, L_2)(L_3) &= \gamma(L_1, L_2, L_3) \end{aligned}$$

for all $L, L_1, L_2, L_3 \in \mathfrak{l}$ and $A, A_1, A_2 \in \mathfrak{a}$. Using that α is a cocycle satisfying (2) it is not hard to prove that β is also a cocycle, i.e. $\beta \in Z^2(\mathfrak{d}_0, \mathfrak{l}^*)$. Then the bilinear map $[\cdot, \cdot]$ is exactly the Lie bracket on $\mathfrak{l}^* \oplus \mathfrak{a} \oplus \mathfrak{l}$ defined by the central extension

$$0 \longrightarrow \mathfrak{l}^* \longrightarrow \mathfrak{l}^* \oplus \mathfrak{a} \oplus \mathfrak{l} \longrightarrow \mathfrak{d}_0 \longrightarrow 0$$

associated with $\beta \in Z^2(\mathfrak{d}_0, \mathfrak{l}^*)$. The invariance of $\langle \cdot, \cdot \rangle$ follows from the orthogonality of ρ . $\square$

By construction, $\mathfrak{l}^*$ is an isotropic subspace of the centre $\mathfrak{z}(\mathfrak{d})$ of $\mathfrak{d} := \mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$.

Definition 2.2 We call $\mathfrak{d} = \mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ regular if and only if $\mathfrak{z}(\mathfrak{d}) = \mathfrak{l}^*$.

Lemma 2.1 $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is not regular if and only if there exist vectors $L_0 \in \mathfrak{l}$ and $A_0 \in \mathfrak{a}$, $L_0 + A_0 \neq 0$, such that

$$\begin{aligned} \rho(L_0) &= 0 \\ \rho(\cdot)(A_0) &= \alpha(L_0, \cdot) \\ \gamma(L_0, \cdot, \cdot) &= \langle A_0, \alpha(\cdot, \cdot) \rangle. \end{aligned} \tag{3}$$

In other words, $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is not regular if and only if Equations (3) are satisfied for some L_0, A_0 with $L_0 \neq 0$ or the subspace $\mathfrak{a}^\perp \cap \alpha(\mathfrak{l}, \mathfrak{l})^\perp \subset \mathfrak{a}$ is non-zero. Here $\mathfrak{a}^\perp$ denotes the space of invariants of ρ .

Proof. A straightforward computation shows that an element $X = L_0 + A_0 + Z_0 \in \mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho) = \mathfrak{l} \oplus \mathfrak{a} \oplus \mathfrak{l}^*$ is central if and only if the Equations (3) are satisfied. $\square$

For any metric Lie algebra $\mathfrak{g}$ we have that $\mathfrak{z}(\mathfrak{g})^\perp = \mathfrak{g}'$. Thus $\mathfrak{d} = \mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is regular if and only if $\mathfrak{d}' = \mathfrak{l}^* \oplus \mathfrak{a}$. In particular, in the regular case $\mathfrak{z}(\mathfrak{d}) \subset \mathfrak{d}'$, and $\mathfrak{d}'/\mathfrak{z}(\mathfrak{d})$ is abelian. Now we will prove a converse statement. Note that the centre of a non-abelian indecomposable metric Lie algebra $\mathfrak{g}$ is isotropic, i.e. $\mathfrak{z}(\mathfrak{g}) \subset \mathfrak{g}'$.

Proposition 2.2 If $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is a non-abelian indecomposable metric Lie algebra such that

$$\mathfrak{g}'/\mathfrak{z}(\mathfrak{g}) \text{ is abelian,} \tag{4}$$

then there exist an abelian Lie algebra $\mathfrak{l}$, a semi-Euclidean vector space $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$, an orthogonal representation ρ of $\mathfrak{l}$ on $\mathfrak{a}$, a cocycle $\alpha \in Z^2(\mathfrak{l}, \mathfrak{a})$ satisfying (2), and a 3-form $\gamma \in \wedge^3 \mathfrak{l}^*$ such that $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is isomorphic to $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$.

In particular, we can choose

$$\begin{aligned} \mathfrak{a} &= \mathfrak{g}'/\mathfrak{z}(\mathfrak{g}) \\ \mathfrak{l} &= \mathfrak{g}/\mathfrak{g}' \end{aligned}$$

and

$$\rho(L)(A) = [l, a] + \mathfrak{z}(\mathfrak{g})$$

for $l \in \mathfrak{g}$, $a \in \mathfrak{g}'$, $L = l + \mathfrak{g}'$, $A = a + \mathfrak{z}(\mathfrak{g})$. With this choice, $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is regular.

Proof. We first give the idea of the proof. Choose $\mathfrak{a}$, $\mathfrak{l}$ and ρ as above. Note, that because of (4) ρ is correctly defined. We abbreviate the notation by $\mathfrak{z} = \mathfrak{z}(\mathfrak{g})$. By assumption the Lie algebra $\mathfrak{g}$ is the result of the following two extensions with abelian kernel:

$$\begin{aligned} 0 &\longrightarrow \mathfrak{a} \longrightarrow \mathfrak{g}/\mathfrak{g}' \longrightarrow \mathfrak{l} \longrightarrow 0 \\ 0 &\longrightarrow \mathfrak{z} \longrightarrow \mathfrak{g} \longrightarrow \mathfrak{g}/\mathfrak{z} \longrightarrow 0. \end{aligned}$$

The first extension is given by the representation ρ as defined above and, using an appropriate split, by a cocycle $\alpha \in Z^2(\mathfrak{l}, \mathfrak{a})$. The cocycle defining the second extension will determine $\gamma \in \bigwedge^3 \mathfrak{l}^*$.

Since $\mathfrak{g}' = \mathfrak{z}^\perp$ we have $\mathfrak{a} = \mathfrak{z}^\perp/\mathfrak{z}$. Therefore $\langle \cdot, \cdot \rangle$ induces a non-degenerate inner product $\langle \cdot, \cdot \rangle_{\mathfrak{a}}$ on $\mathfrak{a}$. Because of the ad-invariance of $\langle \cdot, \cdot \rangle$ the representation ρ is orthogonal with respect to $\langle \cdot, \cdot \rangle_{\mathfrak{a}}$.

Because of $\mathfrak{g}' = \mathfrak{z}^\perp$ we can choose an isotropic complementary subspace of $\mathfrak{g}'$ in $\mathfrak{g}$ and identify it with $\mathfrak{l}$ (as a vector space). This yields a section

$$s : \mathfrak{l} \longrightarrow \mathfrak{g}.$$

Since $s(\mathfrak{l})$ is an isotropic complement of $\mathfrak{g}' = \mathfrak{z}^\perp$ in $\mathfrak{g}$ the inner product $\langle \cdot, \cdot \rangle$ defines a dual pairing between $s(\mathfrak{l})$ and $\mathfrak{z}$. Therefore $s(\mathfrak{l})^\perp \cap \mathfrak{g}'$ is a (non-degenerate) complement of $\mathfrak{z}$ in $\mathfrak{g}'$. Identifying $\mathfrak{a}$ with $s(\mathfrak{l})^\perp \cap \mathfrak{g}'$ we obtain a section

$$t : \mathfrak{a} \longrightarrow \mathfrak{g}.$$

We define a cocycle $\alpha \in Z^2(\mathfrak{l}, \mathfrak{a})$ by

$$\alpha(L_1, L_2) = [s(L_1), s(L_2)] + \mathfrak{z}.$$

Then α satisfies Equation (2). This follows from

$$\begin{aligned} \langle \alpha(L_1, L_2), \alpha(L_3, L_4) \rangle_{\mathfrak{a}} &= \langle [s(L_1), s(L_2)], [s(L_3), s(L_4)] \rangle \\ &= \langle [[s(L_1), s(L_2)], s(L_3)], s(L_4) \rangle \end{aligned}$$

and the Jacobi identity in $\mathfrak{g}$. Finally we define a 3-form $\gamma \in \bigwedge^3 \mathfrak{l}^*$ by

$$\gamma(L_1, L_2, L_3) = \langle [s(L_1), s(L_2)], s(L_3) \rangle.$$

Obviously we have an isomorphism

$$i : \mathfrak{l}^* \longrightarrow \mathfrak{z}$$

by $\langle i(Z), s(L) \rangle = Z(L)$ for $Z \in \mathfrak{l}^*$ and $L \in \mathfrak{l}$. Now we will prove that the vector space isomorphism

$$i + t + s : \mathfrak{l}^* \oplus \mathfrak{a} \oplus \mathfrak{l} \longrightarrow \mathfrak{g}$$

is an isomorphism between the metric Lie algebras $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ and $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$. It is almost obvious that the map $i + t + s$ is an isometry. Thus it remains to show that it is an

isomorphism of Lie algebras. This follows from the following equations which use the invariance of $\langle \cdot, \cdot \rangle$:

$$\begin{aligned}
[A_1, A_2] &= \langle \rho(\cdot)A_1, A_2 \rangle_{\mathfrak{a}} \\
[t(A_1), t(A_2)] &\in \mathfrak{z} \quad \text{by (4)} \\
\langle [t(A_1), t(A_2)], s(L) \rangle &= \langle [s(L), t(A_1)], t(A_2) \rangle = \langle \rho(L)(A_1), A_2 \rangle_{\mathfrak{a}} \\
[L, A] &= -\langle A, \alpha(L, \cdot) \rangle_{\mathfrak{a}} + \rho(L)(A) \\
[s(L), t(A)] &\in \mathfrak{g}' \\
[s(L), t(A)] &\equiv t(\rho(L)(A)) \pmod{\mathfrak{z}} \\
\langle [s(L), t(A)], s(L_1) \rangle_{\mathfrak{a}} &= -\langle t(A), [s(L), s(L_1)] \rangle = -\langle A, \alpha(L, L_1) \rangle_{\mathfrak{a}} \\
[L_1, L_2] &= \gamma(L_1, L_2, \cdot) + \alpha(L_1, L_2) \\
[s(L_1), s(L_2)] &\in \mathfrak{g}' \\
[s(L_1), s(L_2)] &\equiv t(\alpha(L_1, L_2)) \pmod{\mathfrak{z}} \\
\langle [s(L_1), s(L_2)], s(L_3) \rangle &= \gamma(L_1, L_2, L_3).
\end{aligned}$$

□

Remark 2.1 It is natural to ask to what extent the tuple (α, γ) is determined by $\mathfrak{g}$. The above proof gives that (α, γ) depends on the choice of a section $s : \mathfrak{l} \rightarrow \mathfrak{g}$. A closer view shows that (α, γ) depends on the induced section $\bar{s} : \mathfrak{l} \rightarrow \mathfrak{g}/\mathfrak{z}$, only. This corresponds to the fact that the orbit of (α, γ) under a certain action of the group of 1-cocycles $C^1(\mathfrak{l}, \mathfrak{a})$ is canonically attached to $\mathfrak{g}$ (see Section 3).

Definition 2.3 Let $\mathfrak{i}$ be an isotropic ideal in $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$. We will say that

1. $\mathfrak{i}$ is maximal isotropic if $\dim \mathfrak{i} = \min(p, q)$, where (p, q) is the signature of $\langle \cdot, \cdot \rangle$,
2. $\mathfrak{i}$ is isomaximal if $\mathfrak{i}$ is not properly contained in a further isotropic ideal.

Obviously, each maximal isotropic ideal is isomaximal.

Proposition 2.3 Let $\mathfrak{g}$ be an indecomposable metric Lie algebra with non-trivial isomaximal centre. Then $\mathfrak{g}$ satisfies condition (4).

Proof. We have to show that $\mathfrak{z}(\mathfrak{g})^\perp/\mathfrak{z}(\mathfrak{g})$ is abelian. In Lemma 2.3 we will prove even more, namely that $\mathfrak{i}^\perp/\mathfrak{i}$ is abelian for each isomaximal ideal $\mathfrak{i}$. □

Lemma 2.2 Let $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ be an indecomposable metric Lie algebra and $\mathfrak{r}$ the radical of $\mathfrak{g}$. Assume that $\mathfrak{r} \neq 0$. Then $\mathfrak{r}^\perp \subset \mathfrak{r}$ holds.

Proof. Let $\mathfrak{g} = \mathfrak{s} \ltimes \mathfrak{r}$ be a Levi decomposition of $\mathfrak{g}$ and consider $\mathfrak{s}_1 := \mathfrak{s} \cap \mathfrak{r}^\perp$. Clearly, $\mathfrak{s}_1$ is an ideal in $\mathfrak{s}$. Since $\mathfrak{s}$ is semi-simple, there is a complementary ideal $\mathfrak{s}_2$ of $\mathfrak{s}_1$ in $\mathfrak{s}$, i.e. $\mathfrak{s} = \mathfrak{s}_1 \oplus \mathfrak{s}_2$ as a direct sum of ideals. It is easy to see that $\mathfrak{s}_2 \ltimes \mathfrak{r}$ is a non-degenerate ideal in $\mathfrak{g}$. Since $\mathfrak{g}$ is indecomposable and $\mathfrak{r} \neq 0$ we obtain $\mathfrak{s}_2 \ltimes \mathfrak{r} = \mathfrak{g}$. Hence, $\mathfrak{s}_1 = 0$ and, consequently, $\mathfrak{r}^\perp \subset \mathfrak{r}$. □

Lemma 2.3 *Let $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ be a metric Lie algebra with non-trivial radical $\mathfrak{r} \neq 0$ and let $\mathfrak{i} \subset \mathfrak{g}$ be an isomaximal ideal. Then we have*

1. $\mathfrak{r} \cap \mathfrak{r}^\perp \subset \mathfrak{i}$.

If, in addition, $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is indecomposable, then

2. $\mathfrak{i}^\perp \subset \mathfrak{r}$,

3. $\mathfrak{i}^\perp/\mathfrak{i}$ is abelian.

Proof. Since the ideal $\mathfrak{i}$ is isotropic it is abelian. Hence, it is contained in the radical $\mathfrak{r}$. In particular, $\langle \mathfrak{i}, \mathfrak{r} \cap \mathfrak{r}^\perp \rangle = 0$. Therefore, $\mathfrak{i} + \mathfrak{r} \cap \mathfrak{r}^\perp$ is an isotropic ideal, which must be equal to $\mathfrak{i}$ since $\mathfrak{i}$ is isomaximal. This implies $\mathfrak{r} \cap \mathfrak{r}^\perp \subset \mathfrak{i}$. Using this we obtain

$$\mathfrak{i}^\perp \subset (\mathfrak{r} \cap \mathfrak{r}^\perp)^\perp = \mathfrak{r}^\perp + \mathfrak{r} \subset \mathfrak{r},$$

where the last inclusion uses Lemma 2.2. Now we are going to prove the third assertion. Since $\mathfrak{i}^\perp \subset \mathfrak{r}$ the ideal $\mathfrak{i}^\perp$ is solvable. Therefore, also

$$\mathfrak{a} := \mathfrak{i}^\perp/\mathfrak{i}$$

is solvable. We define an ideal $\tilde{\mathfrak{z}}$ of $\mathfrak{i}^\perp$ by

$$\mathfrak{i} \subset \tilde{\mathfrak{z}} := \{X \in \mathfrak{i}^\perp \mid [X, Y] \in \mathfrak{i} \text{ for all } Y \in \mathfrak{i}^\perp\} \subset \mathfrak{i}^\perp.$$

Obviously, $\tilde{\mathfrak{z}}/\mathfrak{i}$ is the centre of $\mathfrak{a}$. Consider now

$$\mathfrak{j} := [\mathfrak{i}^\perp, \mathfrak{i}^\perp] \cap \tilde{\mathfrak{z}} + \mathfrak{i}.$$

Since $\mathfrak{i}, [\mathfrak{i}^\perp, \mathfrak{i}^\perp]$ and $\tilde{\mathfrak{z}}$ are ideals $\mathfrak{j}$ is also an ideal. We claim that $\mathfrak{j}$ is isotropic. Indeed, $[\mathfrak{i}^\perp, \mathfrak{i}^\perp] \cap \tilde{\mathfrak{z}}$ is isotropic, because of $[\mathfrak{i}^\perp, \mathfrak{i}^\perp] \perp \tilde{\mathfrak{z}}$. On the other hand, $\mathfrak{i}$ is isotropic and $[\mathfrak{i}^\perp, \mathfrak{i}^\perp] \cap \tilde{\mathfrak{z}}$ is contained in $\mathfrak{i}^\perp$. Thus, $\mathfrak{j}$ is an isotropic ideal containing $\mathfrak{i}$. Therefore, $\mathfrak{j} = \mathfrak{i}$, which implies $[\mathfrak{i}^\perp, \mathfrak{i}^\perp] \cap \tilde{\mathfrak{z}} \subset \mathfrak{i}$. Factorizing this by $\mathfrak{i}$ yields

$$[\mathfrak{a}, \mathfrak{a}] \cap \mathfrak{z}(\mathfrak{a}) = 0, \tag{5}$$

where $\mathfrak{z}(\mathfrak{a})$ denotes the centre of $\mathfrak{a}$. On the other hand we have

$$[\mathfrak{a}, \mathfrak{a}]^\perp = \mathfrak{z}(\mathfrak{a}), \tag{6}$$

Since $\langle \cdot, \cdot \rangle$ induces a non-degenerate ad-invariant inner product on $\mathfrak{a}$. Combining Equations (5) and (6) we obtain

$$\mathfrak{a} = \mathfrak{a}' \oplus \mathfrak{z}(\mathfrak{a}),$$

which implies

$$\mathfrak{a}' = [\mathfrak{a}', \mathfrak{a}'] = \mathfrak{a}^{(2)}.$$

Now we use that $\mathfrak{a}$ is solvable and obtain $\mathfrak{a}' = 0$, which proves the assertion. $\square$

In particular, we can apply Propositions 2.2 and 2.3 to indecomposable metric Lie algebras with maximal isotropic centre $\mathfrak{z}$. In this case the induced inner product on $\mathfrak{z}^\perp/\mathfrak{z}$ is definite. We may assume, perhaps after a sign change of $\langle \cdot, \cdot \rangle$, that it is Euclidean. We obtain the following structure theorem.

Theorem 2.1 *If $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is an indecomposable metric Lie algebra of signature (p, q) , $p \leq q$, with non-trivial maximal isotropic centre, then there exist an abelian Lie algebra $\mathfrak{l}$, a Euclidean vector space $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$, an orthogonal representation ρ of $\mathfrak{l}$ on $\mathfrak{a}$, a cocycle $\alpha \in Z^2(\mathfrak{l}, \mathfrak{a})$ satisfying (2), and a 3-form $\gamma \in \wedge^3 \mathfrak{l}^*$ such that $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is regular and $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is isomorphic to $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$.*

Proposition 2.4 *Let $\mathfrak{g}$ be a non-simple indecomposable metric Lie algebra of index at most 2. Then $\mathfrak{g}$ satisfies condition (4).*

Lemma 2.4 *Let $\mathfrak{g}$ be as in Proposition 2.4. Then $\mathfrak{g}$ is solvable.*

Proof. First we observe that $\mathfrak{g}$ cannot be semi-simple. Thus $\mathfrak{g}$ has a Levi decomposition $\mathfrak{g} = \mathfrak{s} \ltimes \mathfrak{r}$ with $\mathfrak{r} \neq 0$. By Lemma 2.2 we have $\mathfrak{r}^{\perp} \subset \mathfrak{r}$. Therefore the scalar product on $\mathfrak{g}$ induces a non-degenerate pairing between $\mathfrak{s}$ and $\mathfrak{r}^{\perp}$. It follows that the index of $\mathfrak{g}$ is greater or equal to $\dim \mathfrak{s} = \dim \mathfrak{r}^{\perp}$ which is at least 3 if not zero. We conclude that $\mathfrak{s} = 0$. $\square$

Lemma 2.5 *A nilpotent metric Lie algebra of index at most 1 is abelian.*

Proof. This is known according to [M 85], see also [BK 02]. For the convenience of the reader we give a selfcontained argument in the language developed in the present paper. Let $\mathfrak{n}$ be a nilpotent metric Lie algebra of index at most 1. Then $\mathfrak{n}$ is the direct sum of a Euclidean abelian Lie algebra $\mathfrak{n}_0$ and, possibly, an indecomposable nilpotent Lie algebra $\mathfrak{n}_1$ of index 1. We have to show that $\mathfrak{n}_1$ is abelian. Let $\mathfrak{z}$ be the centre of $\mathfrak{n}_1$. It is one-dimensional and maximal isotropic in $\mathfrak{n}_1$. By Theorem 2.1 we have $\mathfrak{n}_1 \cong \mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ with $\mathfrak{l}$ one-dimensional, $\alpha = \gamma = 0$, and $\mathfrak{a} \cong \mathfrak{n}'_1/\mathfrak{z}$ Euclidean. Let H be a generator of $\mathfrak{l}$. Since $\mathfrak{n}_1$ is nilpotent the anti-symmetric endomorphism $\rho(H)$ has to be nilpotent. It follows that $\rho(H) = 0$. Therefore $\mathfrak{n}_1 \cong \mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is abelian. $\square$

Proof of Proposition 2.4. According to Lemma 2.4 the Lie algebra $\mathfrak{g}$ is solvable. By Lie's Theorem $\mathfrak{g}'$ is nilpotent. Thus $\mathfrak{g}'/\mathfrak{z}$ is a nilpotent metric Lie algebra of index at most 1. It is abelian by Lemma 2.5. $\square$

Combining Proposition 2.4 with Proposition 2.2 we obtain

Corollary 2.1 *If $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is a non-simple indecomposable metric Lie algebra of index at most 2, then there exist an abelian Lie algebra $\mathfrak{l}$, $\dim \mathfrak{l} = \dim \mathfrak{z}$, a semi-Euclidean vector space $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$, an orthogonal representation ρ of $\mathfrak{l}$ on $\mathfrak{a}$, and a cocycle $\alpha \in Z^2(\mathfrak{l}, \mathfrak{a})$ satisfying (2), such that $\mathfrak{d}_{\alpha, 0}(\mathfrak{a}, \mathfrak{l}, \rho)$ is regular and $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is isomorphic to $\mathfrak{d}_{\alpha, 0}(\mathfrak{a}, \mathfrak{l}, \rho)$.*

3 Equivalence of twofold extensions and the isomorphism problem

In order to approach the question whether two metric Lie algebras $\mathfrak{d}_{\alpha_i, \gamma_i}(\mathfrak{a}_i, \mathfrak{l}_i, \rho_i)$, $i = 1, 2$, are isomorphic we first introduce a stronger equivalence relation on these objects.

We consider extensions of a Lie algebra $\mathfrak{h}$ by an abelian Lie algebra $\mathfrak{a}$

$$0 \rightarrow \mathfrak{a} \xrightarrow{i} \mathfrak{g} \xrightarrow{p} \mathfrak{h} \rightarrow 0 .$$

If $\mathfrak{h}$ and $\mathfrak{a}$ are understood we denote such an extension by $(\mathfrak{g}, i, p)$. Two such extensions $(\mathfrak{g}_j, i_j, p_j)$, $j = 1, 2$, are called to be equivalent if and only if there exists a Lie algebra isomorphism $\Phi : \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ such that $\Phi \circ i_1 = i_2$ and $p_2 \circ \Phi = p_1$. It is well-known that $(\mathfrak{g}_1, i_1, p_1) \cong (\mathfrak{g}_2, i_2, p_2)$ if and only if the corresponding representations of $\mathfrak{h}$ on $\mathfrak{a}$ coincide and the cocycles $\alpha_j \in Z^2(\mathfrak{h}, \mathfrak{a})$ defined after a choice of sections of p_j are cohomologous. In particular, equivalence classes of extensions with a fixed representation of $\mathfrak{h}$ on $\mathfrak{a}$ are in bijective correspondence with elements of $H^2(\mathfrak{h}, \mathfrak{a})$.

Recall from the proof of Proposition 2.1 that $\mathfrak{d}_{\alpha, \gamma} := \mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is the result of two subsequent extensions of Lie algebras

$$\begin{aligned} (\mathfrak{d}_{\alpha}, i, p) : \quad & 0 \rightarrow \mathfrak{a} \xrightarrow{i} \mathfrak{d}_{\alpha} \xrightarrow{p} \mathfrak{l} \rightarrow 0 , \\ (\mathfrak{d}_{\alpha, \gamma}, j, q) : \quad & 0 \rightarrow \mathfrak{l}^* \xrightarrow{j} \mathfrak{d}_{\alpha, \gamma} \xrightarrow{q} \mathfrak{d}_{\alpha} \rightarrow 0 , \end{aligned}$$

where $\mathfrak{l}$ is abelian, and the representations of $\mathfrak{l}$ on $\mathfrak{a}$ and of $\mathfrak{d}_{\alpha}$ on $\mathfrak{l}^*$ are orthogonal and trivial, respectively.

Definition 3.1 Let $\alpha_i \in Z^2(\mathfrak{l}, \mathfrak{a})$, $i = 1, 2$, be cocycles satisfying (2), and let $\gamma_i \in \bigwedge^3 \mathfrak{l}^*$. We call the metric Lie algebras $\mathfrak{d}_{\alpha_i, \gamma_i}(\mathfrak{a}, \mathfrak{l}, \rho)$ extension equivalent if and only if there exist an equivalence of extensions

$$\Phi : (\mathfrak{d}_{\alpha_1}, i_1, p_1) \rightarrow (\mathfrak{d}_{\alpha_2}, i_2, p_2)$$

and an isomorphism of metric Lie algebras

$$\Psi : \mathfrak{d}_{\alpha_1, \gamma_1}(\mathfrak{a}, \mathfrak{l}, \rho) \rightarrow \mathfrak{d}_{\alpha_2, \gamma_2}(\mathfrak{a}, \mathfrak{l}, \rho)$$

making the following diagram commutative

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathfrak{l}^* & \xrightarrow{j_1} & \mathfrak{d}_{\alpha_1, \gamma_1} & \xrightarrow{q_1} & \mathfrak{d}_{\alpha_1} \longrightarrow 0 \\ & & \downarrow \text{Id} & & \downarrow \Psi & & \downarrow \Phi \\ 0 & \longrightarrow & \mathfrak{l}^* & \xrightarrow{j_2} & \mathfrak{d}_{\alpha_2, \gamma_2} & \xrightarrow{q_2} & \mathfrak{d}_{\alpha_2} \longrightarrow 0 \end{array} . \quad (7)$$

We now consider the standard Lie algebra cochain complex of the abelian Lie algebra $\mathfrak{l}$ with values in $\mathfrak{a}$

$$0 \rightarrow C^0(\mathfrak{l}, \mathfrak{a}) \xrightarrow{d} C^1(\mathfrak{l}, \mathfrak{a}) \xrightarrow{d} C^2(\mathfrak{l}, \mathfrak{a}) \xrightarrow{d} \dots \xrightarrow{d} C^l(\mathfrak{l}, \mathfrak{a}) \rightarrow 0 ,$$

where $C^p(\mathfrak{l}, \mathfrak{a}) = \text{Hom}(\bigwedge^p \mathfrak{l}, \mathfrak{a})$, $l = \dim \mathfrak{l}$, and for $\tau \in C^p(\mathfrak{l}, \mathfrak{a})$

$$d\tau(L_1, \dots, L_{p+1}) = \sum_{i=1}^{p+1} (-1)^{i-1} \rho(L_i) \tau(L_1, \dots, \hat{L}_i, \dots, L_{p+1}) .$$

The composition of maps

$$C^p(\mathfrak{l}, \mathfrak{a}) \times C^q(\mathfrak{l}, \mathfrak{a}) \xrightarrow{\wedge} C^{p+q}(\mathfrak{l}, \mathfrak{a} \otimes \mathfrak{a}) \xrightarrow{\langle \cdot, \cdot \rangle_{\mathfrak{a}}} \wedge^{p+q} \mathfrak{l}^*$$

defines a bilinear multiplication

$$C^p(\mathfrak{l}, \mathfrak{a}) \times C^q(\mathfrak{l}, \mathfrak{a}) \rightarrow \wedge^{p+q} \mathfrak{l}^*$$

which we will denote by $(\alpha, \tau) \mapsto \langle \alpha \wedge \tau \rangle_{\mathfrak{a}}$. We are particularly interested in the case $p = 2, q = 1$. Then

$$\langle \alpha \wedge \tau \rangle_{\mathfrak{a}}(L_1, L_2, L_3) = \langle \alpha(L_1, L_2), \tau(L_3) \rangle_{\mathfrak{a}} + \langle \alpha(L_3, L_1), \tau(L_2) \rangle_{\mathfrak{a}} + \langle \alpha(L_2, L_3), \tau(L_1) \rangle_{\mathfrak{a}} .$$

Proposition 3.1 *The metric Lie algebras $\mathfrak{d}_{\alpha_i, \gamma_i}(\mathfrak{a}, \mathfrak{l}, \rho)$ are extension equivalent if and only if there exists a cochain $\tau \in C^1(\mathfrak{l}, \mathfrak{a})$ such that*

$$\alpha_2 - \alpha_1 = d\tau \tag{8}$$

and

$$\gamma_2 - \gamma_1 = \langle (\alpha_1 + \frac{1}{2}d\tau) \wedge \tau \rangle_{\mathfrak{a}} . \tag{9}$$

Proof. Assume that $\mathfrak{d}_{\alpha_i, \gamma_i}(\mathfrak{a}, \mathfrak{l}, \rho)$ are extension equivalent, i.e., there exist isomorphisms Φ and Ψ as in Definition 3.1. We use the vector space decomposition $\mathfrak{d}_{\alpha_1} = \mathfrak{l} \oplus \mathfrak{a} = \mathfrak{d}_{\alpha_2}$ in order to define $\tau \in C^1(\mathfrak{l}, \mathfrak{a})$ by $\tau(L) := L - \Phi(L)$, $L \in \mathfrak{l}$. Because of $p_2(L - \Phi(L)) = L - p_1(L) = 0$ we have $\tau(L) \in \mathfrak{a}$. Moreover, (8) holds. Indeed,

$$\Phi^{-1}(L) = L + \tau(L)$$

and

$$\begin{aligned} \alpha_2(L_1, L_2) = [L_1, L_2]_{\mathfrak{d}_{\alpha_2}} &= [\Phi^{-1}(L_1), \Phi^{-1}(L_2)]_{\mathfrak{d}_{\alpha_1}} \\ &= [L_1 + \tau(L_1), L_2 + \tau(L_2)]_{\mathfrak{d}_{\alpha_1}} = \alpha_1(L_1, L_2) + d\tau(L_1, L_2) . \end{aligned}$$

We denote the commutator in $\mathfrak{d}_{\alpha_i, \gamma_i}(\mathfrak{a}, \mathfrak{l}, \rho)$ by $[\cdot, \cdot]_i$. The decomposition $\mathfrak{d}_{\alpha_1, \gamma_1}(\mathfrak{a}, \mathfrak{l}, \rho) = \mathfrak{l} \oplus \mathfrak{a} \oplus \mathfrak{l}^* = \mathfrak{d}_{\alpha_2, \gamma_2}(\mathfrak{a}, \mathfrak{l}, \rho)$ will us allow to consider $\mathfrak{l} \oplus \mathfrak{a}$ as a subspace of $\mathfrak{d}_{\alpha_1, \gamma_1}(\mathfrak{a}, \mathfrak{l}, \rho)$ and of $\mathfrak{d}_{\alpha_2, \gamma_2}(\mathfrak{a}, \mathfrak{l}, \rho)$, Φ^{-1} as a map between $\mathfrak{l} \oplus \mathfrak{a}$ and $\mathfrak{d}_{\alpha_1, \gamma_1}(\mathfrak{a}, \mathfrak{l}, \rho)$, etc. We compute

$$\begin{aligned} \gamma_2(L_1, L_2, L) &= \langle [L_1, L_2]_2, L \rangle \\ &= \langle [\Psi^{-1}(L_1), \Psi^{-1}(L_2)]_1, \Psi^{-1}(L) \rangle \\ &= \langle [\Phi^{-1}(L_1), \Phi^{-1}(L_2)]_1, \Phi^{-1}(L) \rangle \\ &= \langle [L_1 + \tau(L_1), L_2 + \tau(L_2)]_1, L + \tau(L) \rangle \\ &= \langle [L_1, L_2]_1, L \rangle \\ &\quad + \langle [L_1, L_2]_1, \tau(L) \rangle + \langle [L_1, \tau(L_2)]_1, L \rangle + \langle [\tau(L_1), L_2]_1, L \rangle \\ &\quad + \langle [\tau(L_1), L_2]_1, \tau(L) \rangle + \langle [L_1, \tau(L_2)]_1, \tau(L) \rangle + \langle [\tau(L_1), \tau(L_2)]_1, L \rangle \\ &= \gamma_1(L_1, L_2, L) \\ &\quad + \langle \alpha_1(L_1, L_2), \tau(L) \rangle_{\mathfrak{a}} + \langle \alpha_1(L, L_1), \tau(L_2) \rangle_{\mathfrak{a}} + \langle \alpha_1(L_2, L), \tau(L_1) \rangle_{\mathfrak{a}} \\ &\quad + \langle \tau(L_1), L_2 \tau(L) \rangle_{\mathfrak{a}} + \langle \tau(L), L_1 \tau(L_2) \rangle_{\mathfrak{a}} + \langle \tau(L_2), L \tau(L_1) \rangle_{\mathfrak{a}} \\ &= \gamma_1(L_1, L_2, L) + \langle \alpha_1 \wedge \tau \rangle_{\mathfrak{a}}(L_1, L_2, L) + \frac{1}{2} \langle d\tau \wedge \tau \rangle_{\mathfrak{a}}(L_1, L_2, L) . \end{aligned} \tag{10}$$

This proves (9).

Vice versa, let us assume that $\tau \in C^1(\mathfrak{l}, \mathfrak{a})$ satisfying (8) and (9) is given. Let $\tau^* : \mathfrak{a} \rightarrow \mathfrak{l}^*$ be its adjoint, where we have identified $\mathfrak{a}$ with $\mathfrak{a}^*$ using $\langle \cdot, \cdot \rangle_{\mathfrak{a}}$. With respect to the decompositions $\mathfrak{d}_{\alpha_i} = \mathfrak{l} \oplus \mathfrak{a}$, $\mathfrak{d}_{\alpha_i, \gamma_i} = \mathfrak{l} \oplus \mathfrak{a} \oplus \mathfrak{l}^*$ we define

$$\begin{aligned}\Phi(L + A) &:= L - \tau(L) + A \\ \Psi(L + A + Z) &:= \Phi(L + A) - \frac{1}{2}\tau^*\tau(L) + \tau^*(A) + Z.\end{aligned}\quad (11)$$

Then we have

$$\Psi = \begin{pmatrix} \text{Id} & \tau^* & -\frac{1}{2}\tau^*\tau \\ 0 & \text{Id} & -\tau \\ 0 & 0 & \text{Id} \end{pmatrix} \quad (12)$$

with respect to the decomposition $\mathfrak{l}^* \oplus \mathfrak{a} \oplus \mathfrak{l}$. Because of (8) the map Φ defines an equivalence between $(\mathfrak{d}_{\alpha_1}, i_1, p_1)$ and $(\mathfrak{d}_{\alpha_2}, i_2, p_2)$. Obviously, Ψ satisfies the relations (7). Moreover, it is easy to verify that Ψ is an isometry.

It remains to show that Ψ is in fact a Lie algebra homomorphism. Using that Ψ is an isometry this is equivalent to

$$\langle [\Psi^{-1}(X_1), \Psi^{-1}(X_2)]_1, \Psi^{-1}(X) \rangle = \langle [X_1, X_2]_2, X \rangle \quad (13)$$

for all $X_i = L_i + A_i + Z_i$, $X = L + A + Z \in \mathfrak{l} \oplus \mathfrak{a} \oplus \mathfrak{l}^*$. We can assume that $Z_i = 0$ since Z_i is central and fixed by Ψ^{-1} and that $Z = 0$ since $[\Psi^{-1}(X_1), \Psi^{-1}(X_2)]_1$ is orthogonal to $\mathfrak{l}^*$. Observe that $\Psi^{-1} - \Phi^{-1}$ maps $\mathfrak{l} \oplus \mathfrak{a}$ to $\mathfrak{l}^*$. Using again that commutators are orthogonal to $\mathfrak{l}^*$, we obtain

$$\langle [\Psi^{-1}(X_1), \Psi^{-1}(X_2)]_1, \Psi^{-1}(X) \rangle = \langle [\Phi^{-1}(X_1), \Phi^{-1}(X_2)]_1, \Phi^{-1}(X) \rangle.$$

If $X = A \in \mathfrak{a}$, then this is equal to

$$\begin{aligned}\langle [\Phi^{-1}(X_1), \Phi^{-1}(X_2)]_1, A \rangle &= \langle [\Phi^{-1}(X_1), \Phi^{-1}(X_2)]_{\mathfrak{d}_{\alpha_1}}, A \rangle \\ &= \langle \Phi^{-1}([X_1, X_2]_{\mathfrak{d}_{\alpha_2}}), A \rangle = \langle [X_1, X_2]_2, A \rangle,\end{aligned}$$

since Φ^{-1} is a Lie algebra homomorphism which is equal to the identity when restricted to $\mathfrak{a}$.

Now let $X = L \in \mathfrak{l}$. Employing that $\Phi^{-1}(L) = L + \tau(L)$ we compute

$$\langle [\Phi^{-1}(A_1), \Phi^{-1}(A_2)]_1, \Phi^{-1}(L) \rangle = \langle [A_1, A_2]_1, L + \tau(L) \rangle = \langle A_2, LA_1 \rangle_{\mathfrak{a}} = \langle [A_1, A_2]_2, L \rangle$$

and

$$\begin{aligned}\langle [\Phi^{-1}(L_1), \Phi^{-1}(A_2)]_1, \Phi^{-1}(L) \rangle &= \langle [L_1 + \tau(L_1), A_2]_1, L + \tau(L) \rangle \\ &= \langle L_1 A_2, \tau(L) \rangle_{\mathfrak{a}} + \langle \alpha_1(L, L_1), A_2 \rangle_{\mathfrak{a}} - \langle \tau(L_1), LA_2 \rangle_{\mathfrak{a}} \\ &= \langle (\alpha_1 + d\tau)(L, L_1), A_2 \rangle_{\mathfrak{a}} = \langle \alpha_2(L, L_1), A_2 \rangle_{\mathfrak{a}} \\ &= \langle [L_1, A_2]_2, L \rangle.\end{aligned}$$

Eventually we have (repeating the computation (10))

$$\begin{aligned}\langle [\Phi^{-1}(L_1), \Phi^{-1}(L_2)]_1, \Phi^{-1}(L) \rangle &= \langle [L_1 + \tau(L_1), L_2 + \tau(L_2)]_1, L + \tau(L) \rangle \\ &= \gamma_1(L_1, L_2, L) + \langle (\alpha_1 + \frac{1}{2}d\tau) \wedge \tau \rangle_{\mathfrak{a}}(L_1, L_2, L) \\ &= \gamma_2(L_1, L_2, L) = \langle [L_1, L_2]_2, L \rangle.\end{aligned}$$

This finishes the proof of (13) and, hence, of the proposition. $\square$

Now we are going to construct a certain set which should parametrize the extension equivalence classes for fixed $(\mathfrak{a}, \mathfrak{l}, \rho)$ in the same way as $H^2(\mathfrak{l}, \mathfrak{a})$ parametrizes extensions of $\mathfrak{l}$ by the $\mathfrak{l}$ -module $(\rho, \mathfrak{a})$.

Since the representation ρ is orthogonal the product $\langle \cdot \wedge \cdot \rangle_{\mathfrak{a}}$ is compatible with the differential d in the following way

$$\langle d\alpha \wedge \tau \rangle_{\mathfrak{a}} = (-1)^{p+1} \langle \alpha \wedge d\tau \rangle_{\mathfrak{a}}, \quad \alpha \in C^p(\mathfrak{l}, \mathfrak{a}), \tau \in C^q(\mathfrak{l}, \mathfrak{a}). \quad (14)$$

Therefore it induces a kind of cup product on the cohomology groups

$$\cup : H^p(\mathfrak{l}, \mathfrak{a}) \times H^q(\mathfrak{l}, \mathfrak{a}) \rightarrow \bigwedge^{p+q} \mathfrak{l}^*, \quad [\alpha] \cup [\tau] := \langle \alpha \wedge \tau \rangle_{\mathfrak{a}}. \quad (15)$$

Note that the left hand side of (2) is equal to $\frac{1}{2} \langle \alpha \wedge \alpha \rangle_{\mathfrak{a}}$. Thus property (2) only depends on the cohomology class $a := [\alpha] \in H^2(\mathfrak{l}, \mathfrak{a})$ ¹ via the condition

$$a \cup a = 0.$$

Denote by $Z_C^2(\mathfrak{l}, \mathfrak{a}) \subset Z^2(\mathfrak{l}, \mathfrak{a})$ the space of all cocycles satisfying (2).

Since the map $C^1(\mathfrak{l}, \mathfrak{a}) \ni \tau \mapsto \Psi \in GL(\mathfrak{l}^* \oplus \mathfrak{a} \oplus \mathfrak{l})$ given by (12) is a group homomorphism up to antisymmetric maps from $\mathfrak{l}$ to $\mathfrak{l}^*$ we can expect to define an action of the abelian group of 1-cocycles $C^1(\mathfrak{l}, \mathfrak{a})$, which we write as a right action, on the direct product $C^2(\mathfrak{l}, \mathfrak{a}) \times \bigwedge^3 \mathfrak{l}^*$ by

$$(\alpha, \gamma)\tau := (\alpha, \gamma) + \left(d\tau, \langle (\alpha + \frac{1}{2}d\tau) \wedge \tau \rangle_{\mathfrak{a}} \right), \quad \tau \in C^1(\mathfrak{l}, \mathfrak{a}), \alpha \in C^2(\mathfrak{l}, \mathfrak{a}), \gamma \in \bigwedge^3 \mathfrak{l}^*. \quad (16)$$

Let us verify that (16) in fact defines an action. Using (14) we observe that for $\tau_1, \tau_2 \in C^1(\mathfrak{l}, \mathfrak{a})$

$$\langle d\tau_2 \wedge \tau_1 \rangle_{\mathfrak{a}} = \langle \tau_2 \wedge d\tau_1 \rangle_{\mathfrak{a}} = \langle d\tau_1 \wedge \tau_2 \rangle_{\mathfrak{a}},$$

thus

$$\frac{1}{2}(\langle d\tau_1 \wedge \tau_2 \rangle_{\mathfrak{a}} + \langle d\tau_2 \wedge \tau_1 \rangle_{\mathfrak{a}}) = \langle d\tau_1 \wedge \tau_2 \rangle_{\mathfrak{a}}.$$

Therefore we have

$$\begin{aligned} (\alpha, \gamma)(\tau_1 + \tau_2) &= (\alpha, \gamma) + \left(d(\tau_1 + \tau_2), \langle (\alpha + \frac{1}{2}d(\tau_1 + \tau_2)) \wedge (\tau_1 + \tau_2) \rangle_{\mathfrak{a}} \right) \\ &= \left(\alpha + d\tau_1, \gamma + \langle (\alpha + \frac{1}{2}d\tau_1) \wedge \tau_1 \rangle_{\mathfrak{a}} \right) + \left(d\tau_2, \langle (\alpha + d\tau_1 + \frac{1}{2}d\tau_2) \wedge \tau_2 \rangle_{\mathfrak{a}} \right) \\ &= ((\alpha, \gamma)\tau_1)\tau_2. \end{aligned}$$

The subspaces $Z_C^2(\mathfrak{l}, \mathfrak{a}) \times \bigwedge^3 \mathfrak{l}^* \subset Z^2(\mathfrak{l}, \mathfrak{a}) \times \bigwedge^3 \mathfrak{l}^* \subset C^2(\mathfrak{l}, \mathfrak{a}) \times \bigwedge^3 \mathfrak{l}^*$ are invariant under the action of $C^1(\mathfrak{l}, \mathfrak{a})$. The desired parameter set will be the orbit space

$$\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}) := (Z_C^2(\mathfrak{l}, \mathfrak{a}) \times \bigwedge^3 \mathfrak{l}^*) / C^1(\mathfrak{l}, \mathfrak{a}).$$

We remark that the group of coboundaries $B^1(\mathfrak{l}, \mathfrak{a}) \subset C^1(\mathfrak{l}, \mathfrak{a})$ acts trivially on $Z^2(\mathfrak{l}, \mathfrak{a}) \times \bigwedge^3 \mathfrak{l}^*$ because of (14). If $(\alpha, \gamma) \in Z_C^2(\mathfrak{l}, \mathfrak{a}) \times \bigwedge^3 \mathfrak{l}^*$, then we denote by $[\alpha, \gamma]$ the corresponding element in $\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a})$.

Now the following corollary is just a reformulation of Proposition 3.1.

¹This fact is already implicitly contained in Proposition 3.1. Indeed, its proof shows that if (α, γ) defines a metric Lie algebra, then for *any* $\tau \in C^1(\mathfrak{l}, \mathfrak{a})$ the tuple $(\alpha + d\tau, \gamma + \langle (\alpha + \frac{1}{2}d\tau) \wedge \tau \rangle_{\mathfrak{a}})$ defines a metric Lie algebra, too. The Jacobi identity of the latter algebra implies that $\alpha + d\tau$ satisfies (2).

Corollary 3.1 *Fix $(\mathfrak{a}, \mathfrak{l}, \rho)$. The correspondence*

$$\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho) \longmapsto [\alpha, \gamma] \in \mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a})$$

defines a bijection between the extension equivalence classes of metric Lie algebras of the form $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ and elements of $\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a})$.

Remark 3.1 The set $\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a})$ appears among the cohomology sets introduced and studied by Grishkov in [Gr 98]. There it is denoted by H_Δ^3 , where Δ is the multiplication given by $\frac{1}{2}\langle \cdot \wedge \cdot \rangle_{\mathfrak{a}}$. This paper also suggests how one should define $\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a})$ for non-abelian $\mathfrak{l}$ such that the analogue of Corollary 3.1 remains true (compare Prop. 3.2 of [Gr 98] which, however, seems to be not quite correct).

Remark 3.2 Since $\mathfrak{l}$ is abelian the operators $\rho(L)$, $L \in \mathfrak{l}$, have a joint Jordan decomposition which gives rise to an $\mathfrak{l}$ -invariant orthogonal decomposition

$$\mathfrak{a} = \mathfrak{a}^{(\mathfrak{l})} \oplus \mathfrak{a}^R,$$

where $\mathfrak{a}^{(\mathfrak{l})}$ is the nil-subspace of the $\mathfrak{l}$ -action on $\mathfrak{a}$ and $\rho(L)|_{\mathfrak{a}^R}$ is invertible for at least one $L \in \mathfrak{l}$ (in fact for L in an open dense subset). As it is well-known, this implies that for all p

$$H^p(\mathfrak{l}, \mathfrak{a}^R) = \{0\} \quad \text{and therefore} \quad H^p(\mathfrak{l}, \mathfrak{a}^{(\mathfrak{l})}) \cong H^p(\mathfrak{l}, \mathfrak{a}). \quad (17)$$

We claim that the embedding

$$Z_C^2(\mathfrak{l}, \mathfrak{a}^{(\mathfrak{l})}) \times \wedge^3 \mathfrak{l}^* \hookrightarrow Z_C^2(\mathfrak{l}, \mathfrak{a}) \times \wedge^3 \mathfrak{l}^*$$

induces an isomorphism

$$\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}^{(\mathfrak{l})}) \cong \mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}). \quad (18)$$

Indeed, using (17) for $p = 1, 2$ and that $B^1(\mathfrak{l}, \mathfrak{a})$ acts trivially on $Z^2(\mathfrak{l}, \mathfrak{a}) \times \wedge^3 \mathfrak{l}^*$ we obtain

$$\begin{aligned} \mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}) &= (Z_C^2(\mathfrak{l}, \mathfrak{a}) \times \wedge^3 \mathfrak{l}^*) / C^1(\mathfrak{l}, \mathfrak{a}) \\ &\cong (Z_C^2(\mathfrak{l}, \mathfrak{a}^{(\mathfrak{l})}) \times \wedge^3 \mathfrak{l}^*) / (C^1(\mathfrak{l}, \mathfrak{a}^{(\mathfrak{l})}) \oplus Z^1(\mathfrak{l}, \mathfrak{a}^R)) \\ &= (Z_C^2(\mathfrak{l}, \mathfrak{a}^{(\mathfrak{l})}) \times \wedge^3 \mathfrak{l}^*) / (C^1(\mathfrak{l}, \mathfrak{a}^{(\mathfrak{l})}) \oplus B^1(\mathfrak{l}, \mathfrak{a}^R)) \\ &= (Z_C^2(\mathfrak{l}, \mathfrak{a}^{(\mathfrak{l})}) \times \wedge^3 \mathfrak{l}^*) / C^1(\mathfrak{l}, \mathfrak{a}^{(\mathfrak{l})}) \\ &= \mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}^{(\mathfrak{l})}). \end{aligned}$$

In particular, any extension equivalence class has a representative $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ such that $\alpha \in Z_C^2(\mathfrak{l}, \mathfrak{a}^{(\mathfrak{l})})$.

Now we can decide whether two metric Lie algebras $\mathfrak{d}_{\alpha_i, \gamma_i}(\mathfrak{a}_i, \mathfrak{l}_i, \rho_i)$, $i = 1, 2$, are isomorphic provided that they are regular (see Definition 2.2). If $S : \mathfrak{l}_1 \rightarrow \mathfrak{l}_2$ is a linear map and $\alpha \in C^p(\mathfrak{l}_2, \mathfrak{a}_2)$ then we can form the pull back $S^* \alpha \in C^p(\mathfrak{l}_1, \mathfrak{a}_2)$ given by

$$S^* \alpha(L_1, \dots, L_p) = \alpha(S(L_1), \dots, S(L_p)).$$

We also have the pull back $\wedge^p \mathfrak{l}_2^* \ni \gamma \mapsto S^* \gamma \in \wedge^p \mathfrak{l}_1^*$. Moreover, cochains can be composed with linear maps $T : \mathfrak{a}_2 \rightarrow \mathfrak{a}_1$. We denote such a composition by $T\alpha$. If T is an isometry such that $\rho_1(L) = T\rho_2(S(L))T^{-1}$ for all $L \in \mathfrak{l}_1$ and $\alpha \in Z_C^2(\mathfrak{l}_2, \mathfrak{a}_2)$, then $T S^* \alpha \in Z_C^2(\mathfrak{l}_1, \mathfrak{a}_1)$.

Lemma 3.1 *We consider the metric Lie algebras $\mathfrak{d}_{\alpha_i, \gamma_i}(\mathfrak{a}_i, \mathfrak{l}_i, \rho_i)$, $i = 1, 2$. If there exist an isomorphism $S : \mathfrak{l}_1 \rightarrow \mathfrak{l}_2$ and an isometry $U : \mathfrak{a}_1 \rightarrow \mathfrak{a}_2$ such that*

$$(i) \quad U \circ \rho_1(L) \circ U^{-1} = \rho_2(S(L)), \quad L \in \mathfrak{l}_1, \text{ and}$$

$$(ii) \quad \mathfrak{d}_{\alpha_1, \gamma_1}(\mathfrak{a}_1, \mathfrak{l}_1, \rho_1) \text{ and } \mathfrak{d}_{U^{-1}S^*\alpha_2, S^*\gamma_2}(\mathfrak{a}_1, \mathfrak{l}_1, \rho_1) \text{ are extension equivalent,}$$

then $\mathfrak{d}_{\alpha_1, \gamma_1}(\mathfrak{a}_1, \mathfrak{l}_1, \rho_1)$ and $\mathfrak{d}_{\alpha_2, \gamma_2}(\mathfrak{a}_2, \mathfrak{l}_2, \rho_2)$ are isomorphic.

Conversely, if $\mathfrak{d}_{\alpha_i, \gamma_i}(\mathfrak{a}_i, \mathfrak{l}_i, \rho_i)$, $i = 1, 2$, are regular and isomorphic, then there exist an isomorphism $S : \mathfrak{l}_1 \rightarrow \mathfrak{l}_2$ and an isometry $U : \mathfrak{a}_1 \rightarrow \mathfrak{a}_2$ satisfying Conditions (i) and (ii).

Proof. A straightforward computation shows that if S and U satisfy (i), then $S \oplus U \oplus (S^{-1})^*$ defines an isomorphism between the metric Lie algebras $\mathfrak{d}_{U^{-1}S^*\alpha_2, S^*\gamma_2}(\mathfrak{a}_1, \mathfrak{l}_1, \rho_1)$ and $\mathfrak{d}_{\alpha_2, \gamma_2}(\mathfrak{a}_2, \mathfrak{l}_2, \rho_2)$. Thus, if $\Psi : \mathfrak{d}_{\alpha_1, \gamma_1}(\mathfrak{a}_1, \mathfrak{l}_1, \rho_1) \rightarrow \mathfrak{d}_{U^{-1}S^*\alpha_2, S^*\gamma_2}(\mathfrak{a}_1, \mathfrak{l}_1, \rho_1)$ is an extension equivalence, then $F := (S \oplus U \oplus (S^{-1})^*) \circ \Psi$ is an isometric isomorphism between $\mathfrak{d}_{\alpha_1, \gamma_1}(\mathfrak{a}_1, \mathfrak{l}_1, \rho_1)$ and $\mathfrak{d}_{\alpha_2, \gamma_2}(\mathfrak{a}_2, \mathfrak{l}_2, \rho_2)$.

In order to prove the opposite direction we really need the regularity assumption. We abbreviate $\mathfrak{d}_{\alpha_i, \gamma_i}(\mathfrak{a}_i, \mathfrak{l}_i, \rho_i)$ by $\mathfrak{d}_i$. Let $F : \mathfrak{d}_1 \rightarrow \mathfrak{d}_2$ be an isomorphism. Then $F(\mathfrak{z}(\mathfrak{d}_1)) = \mathfrak{z}(\mathfrak{d}_2)$ and $F(\mathfrak{d}_1') = F(\mathfrak{d}_2')$. Thus by regularity $F(\mathfrak{l}_1^*) = \mathfrak{l}_2^*$ and $F(\mathfrak{a}_1 \oplus \mathfrak{l}_1^*) = \mathfrak{a}_2 \oplus \mathfrak{l}_2^*$. Identifying the quotients $\mathfrak{d}_i/(\mathfrak{a}_i \oplus \mathfrak{l}_i^*)$ and $(\mathfrak{a}_i \oplus \mathfrak{l}_i^*)/\mathfrak{l}_i^*$ with $\mathfrak{l}_i$ and $\mathfrak{a}_i$, respectively, F induces a bijection $S : \mathfrak{l}_1 \rightarrow \mathfrak{l}_2$ and an isometry $U : \mathfrak{a}_1 \rightarrow \mathfrak{a}_2$. Note that $F|_{\mathfrak{l}_1^*} = (S^{-1})^*$. We compute for $L \in \mathfrak{l}_1, A \in \mathfrak{a}_2$

$$\begin{aligned} U \circ \rho_1(L) \circ U^{-1}(A) &= F([L, U^{-1}(A)]) \bmod \mathfrak{l}_2^* \\ &= [F(L), F(U^{-1}(A))] \bmod \mathfrak{l}_2^* = [S(L), A] \bmod \mathfrak{l}_2^* \\ &= \rho_2(S(L)) . \end{aligned}$$

Thus S and U satisfy (i) and, as remarked at the beginning of the proof, $S \oplus U \oplus (S^{-1})^*$ is an isomorphism between $\mathfrak{d}_{U^{-1}S^*\alpha_2, S^*\gamma_2}(\mathfrak{a}_1, \mathfrak{l}_1, \rho_1)$ and $\mathfrak{d}_2$. Consequently, $\Psi := (S^{-1} \oplus U^{-1} \oplus S^*) \circ F$ is an isomorphism between $\mathfrak{d}_1$ and $\mathfrak{d}_{U^{-1}S^*\alpha_2, S^*\gamma_2}(\mathfrak{a}_1, \mathfrak{l}_1, \rho_1)$ which, by construction, acts on each of the subquotients $\mathfrak{l}_1$, $\mathfrak{a}_1$, and $\mathfrak{l}_1^*$ as the identity. Therefore Ψ is an extension equivalence. This finishes the proof of the lemma. $\square$

Combining Proposition 3.1 with the Lemma 3.1 and identifying $(\mathfrak{a}_1, \mathfrak{l}_1)$ with $(\mathfrak{a}_2, \mathfrak{l}_2)$ we obtain

Corollary 3.2 *We consider the metric Lie algebras $\mathfrak{d}_{\alpha_i, \gamma_i}(\mathfrak{a}, \mathfrak{l}, \rho_i)$, $i = 1, 2$. If there exist linear maps $S \in GL(\mathfrak{l})$ and $U \in O(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ satisfying*

$$U \circ \rho_1(L) \circ U^{-1} = \rho_2(S(L)) , \quad L \in \mathfrak{l} , \quad (19)$$

and a linear map $\tau : \mathfrak{l} \rightarrow \mathfrak{a}$, i.e. $\tau \in C^1(\mathfrak{l}, \mathfrak{a})$, such that

$$U^{-1} \circ \alpha_2(S(L_1), S(L_2)) - \alpha_1(L_1, L_2) = L_1 \tau(L_2) - L_2 \tau(L_1) \quad (20)$$

$$\begin{aligned} \gamma_2(SL_1, SL_2, SL_3) - \gamma_1(L_1, L_2, L_3) = \\ \langle \alpha_1(L_1, L_2), \tau(L_3) \rangle_{\mathfrak{a}} + \langle \alpha_1(L_3, L_1), \tau(L_2) \rangle_{\mathfrak{a}} + \langle \alpha_1(L_2, L_3), \tau(L_1) \rangle_{\mathfrak{a}} \\ + \langle \tau(L_1), L_2 \tau(L_3) \rangle_{\mathfrak{a}} + \langle \tau(L_3), L_1 \tau(L_2) \rangle_{\mathfrak{a}} + \langle \tau(L_2), L_3 \tau(L_1) \rangle_{\mathfrak{a}} , \end{aligned} \quad (21)$$

then $\mathfrak{d}_{\alpha_1, \gamma_1}(\mathfrak{a}, \mathfrak{l}, \rho_1)$ and $\mathfrak{d}_{\alpha_2, \gamma_2}(\mathfrak{a}, \mathfrak{l}, \rho_2)$ are isomorphic. In these formulas the action of $\mathfrak{l}$ on $\mathfrak{a}$ is given by ρ_1 .

Conversely, if $\mathfrak{d}_{\alpha_i, \gamma_i}(\mathfrak{a}, \mathfrak{l}, \rho_i)$, $i = 1, 2$, are regular and isomorphic, then there exist linear maps $S \in GL(\mathfrak{l})$, $U \in O(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$, and $\tau \in C^1(\mathfrak{l}, \mathfrak{a})$ satisfying Equations (19), (20), and (21).

4 A decomposability criterion

There is a natural notion of a direct sum of twofold extensions. Namely, $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is a direct sum if and only if there are decompositions $\mathfrak{l} = \mathfrak{l}_1 \oplus \mathfrak{l}_2$, $\mathfrak{a} = \mathfrak{a}_1 \oplus \mathfrak{a}_2$, $\mathfrak{a}_1 \perp \mathfrak{a}_2$, such that $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho) = \mathfrak{d}_{\alpha_1, \gamma_1}(\mathfrak{a}_1, \mathfrak{l}_1, \rho_1) \oplus \mathfrak{d}_{\alpha_2, \gamma_2}(\mathfrak{a}_2, \mathfrak{l}_2, \rho_2)$ for appropriate data $(\rho_i, \alpha_i, \gamma_i)$, $i = 1, 2$. A decomposable metric Lie algebra $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ need not to be a non-trivial direct sum of twofold extensions. However, we have

Lemma 4.1 *If $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is extension equivalent to a non-trivial direct sum of twofold extensions, then it is decomposable. Conversely, if $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is regular and decomposable, then it is extension equivalent to a non-trivial direct sum of twofold extensions.*

Proof. If $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is extension equivalent to a non-trivial direct sum of twofold extensions, then it is isomorphic to a direct sum of two nondegenerate ideals, hence decomposable.

Again, for the opposite direction we need the assumption of regularity. We write $\mathfrak{d}$ for $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ and assume that there is a non-trivial orthogonal decomposition into ideals

$$\mathfrak{d} = \mathfrak{d}_1 \oplus \mathfrak{d}_2 .$$

This induces decompositions $\mathfrak{d}' = \mathfrak{d}'_1 \oplus \mathfrak{d}'_2$ and $\mathfrak{z}(\mathfrak{d}) = \mathfrak{z}(\mathfrak{d}_1) \oplus \mathfrak{z}(\mathfrak{d}_2)$. By regularity we obtain

$$\mathfrak{l}^* = \mathfrak{z}(\mathfrak{d}) = \mathfrak{l}_1^* \oplus \mathfrak{l}_2^* ,$$

where $\mathfrak{l}_i^* := \mathfrak{l}^* \cap \mathfrak{d}_i = \mathfrak{z}(\mathfrak{d}_i)$, $i = 1, 2$, and

$$\mathfrak{a} \oplus \mathfrak{l}^* = \mathfrak{d}'_1 \oplus \mathfrak{d}'_2 .$$

Identifying $\mathfrak{a}$ with the quotient $(\mathfrak{a} \oplus \mathfrak{l}^*)/\mathfrak{l}^*$ we obtain an orthogonal decomposition

$$\mathfrak{a} = \mathfrak{a}_1 \oplus \mathfrak{a}_2 ,$$

where $\mathfrak{a}_i := \mathfrak{d}'_i/\mathfrak{l}_i^*$. Note that $\mathfrak{a}_i$, considered as a subspace of $\mathfrak{d}$, need not to be contained in $\mathfrak{d}_i$. Similarly

$$\mathfrak{l} = \mathfrak{d}/\mathfrak{g}' = \mathfrak{l}_1 \oplus \mathfrak{l}_2 ,$$

where $\mathfrak{l}_i := \mathfrak{d}_i/\mathfrak{d}'_i$. Then the inner product of $\mathfrak{d}_i$ identifies $\mathfrak{l}_i^*$ as defined above with the dual of $\mathfrak{l}_i$. Again, $\mathfrak{l}_i$ considered as a subspace of $\mathfrak{d}$ need not to be contained in $\mathfrak{d}_i$. However, we have exact sequences

$$0 \longrightarrow \mathfrak{a}_i \longrightarrow \mathfrak{d}_i/\mathfrak{l}_i^* \longrightarrow \mathfrak{l}_i \longrightarrow 0$$

and

$$0 \longrightarrow \mathfrak{l}_i^* \longrightarrow \mathfrak{d}_i \longrightarrow \mathfrak{d}_i/\mathfrak{l}_i^* \longrightarrow 0 ,$$

which give $\mathfrak{d}_i$ the structure of a twofold extension. Hence, after the choice of appropriate splits, there are isomorphisms

$$F_i : \mathfrak{d}_i \rightarrow \mathfrak{d}_{\alpha_i, \gamma_i}(\mathfrak{a}_i, \mathfrak{l}_i, \rho_i)$$

for certain data $(\rho_i, \alpha_i, \gamma_i)$ (compare the proof of Proposition 2.2). Let $P_i : \mathfrak{d} \rightarrow \mathfrak{d}_i$ be the projections. Then by construction

$$(F_1 \circ P_1) \oplus (F_2 \circ P_2) : \mathfrak{d} \rightarrow \mathfrak{d}_{\alpha_1, \gamma_1}(\mathfrak{a}_1, \mathfrak{l}_1, \rho_1) \oplus \mathfrak{d}_{\alpha_2, \gamma_2}(\mathfrak{a}_2, \mathfrak{l}_2, \rho_2)$$

is an extension equivalence. This finishes the proof of the lemma. $\square$

Proposition 4.1 *We consider the metric Lie algebra $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$. If*

(i) *there are decompositions*

$$\mathfrak{a} = \mathfrak{a}_1 \oplus \mathfrak{a}_2, \quad \mathfrak{l} = \mathfrak{l}_1 \oplus \mathfrak{l}_2$$

satisfying $\mathfrak{a}_1 \perp \mathfrak{a}_2$ and $\mathfrak{l}_i \oplus \mathfrak{a}_i \neq 0$, $i = 1, 2$, and representations ρ_i of $\mathfrak{l}_i$ on $\mathfrak{a}_i$ such that

$$\rho = \rho_1 \oplus \rho_2,$$

(ii) *the cocycle $\alpha \in Z_C^2(\mathfrak{l}, \mathfrak{a})$ satisfies*

$$\alpha(\mathfrak{l}_1 \times \mathfrak{l}_1) \subset \mathfrak{a}_1, \quad \alpha(\mathfrak{l}_2 \times \mathfrak{l}_2) \subset \mathfrak{a}_2,$$

(iii) *there are linear maps $T_1 : \mathfrak{l}_1 \rightarrow \mathfrak{a}_2$ and $T_2 : \mathfrak{l}_2 \rightarrow \mathfrak{a}_1$ such that*

$$\alpha(L_1, L_2) = L_1 T_2 L_2 - L_2 T_1 L_1$$

and

$$\begin{aligned} \gamma(L_1, L'_1, L_2) &= \langle \alpha(L_1, L'_1), T_2(L_2) \rangle_{\mathfrak{a}} + \langle \alpha(L'_1, L_2), T_1(L_1) \rangle_{\mathfrak{a}}, \\ \gamma(L_2, L'_2, L_1) &= \langle \alpha(L_2, L'_2), T_1(L_1) \rangle_{\mathfrak{a}} + \langle \alpha(L'_2, L_1), T_2(L_2) \rangle_{\mathfrak{a}} \end{aligned}$$

for all $L_1, L'_1 \in \mathfrak{l}_1$ and $L_2, L'_2 \in \mathfrak{l}_2$

then $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is decomposable. Conversely, if $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is regular and decomposable, then Conditions (i), (ii) and (iii) hold.

Proof. Let us start the proof with the following remark. If for $\mathfrak{d}_{\alpha_0, \gamma_0}(\mathfrak{a}, \mathfrak{l}, \rho)$ Conditions (i), (ii) and (iii) are satisfied with $T_1 = T_2 = 0$, then we will say that α_0 and γ_0 are diagonal. If the metric Lie algebra $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is extension equivalent to $\mathfrak{d}_{\alpha_0, \gamma_0}(\mathfrak{a}, \mathfrak{l}, \rho)$ for diagonal α_0 and γ_0 , then it is decomposable by Lemma 4.1. Conversely, if $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is regular and decomposable, then it is extension equivalent to $\mathfrak{d}_{\alpha_0, \gamma_0}(\mathfrak{a}, \mathfrak{l}, \rho)$ with diagonal α_0 and γ_0 .

Now let us first assume that $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is regular and decomposable. Hence $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is extension equivalent to a metric Lie algebra $\mathfrak{d}_{\alpha_0, \gamma_0}(\mathfrak{a}, \mathfrak{l}, \rho)$ with diagonal α_0 and γ_0 . By Proposition 3.1 there exists a linear map $\tau : \mathfrak{l} \rightarrow \mathfrak{a}$ such that

$$\alpha = \alpha_0 + d\tau \quad \text{and} \quad \gamma = \gamma_0 + \langle (\alpha_0 + \frac{1}{2}d\tau) \wedge \tau \rangle_{\mathfrak{a}}.$$

We decompose $\tau = \tau_0 + T_1 + T_2$, where $\tau_0(\mathfrak{l}_i) \subset \mathfrak{a}_i$, $T_1(\mathfrak{l}_2) = T_2(\mathfrak{l}_1) = 0$, and $T_1(\mathfrak{l}_1) \subset \mathfrak{a}_2$, $T_2(\mathfrak{l}_2) \subset \mathfrak{a}_1$. Using that $\rho(L_1)|_{\mathfrak{a}_2} = \rho(L_2)|_{\mathfrak{a}_1} = 0$ for $L_i \in \mathfrak{l}_i$ we obtain

$$d\tau(\mathfrak{l}_i \times \mathfrak{l}_i) = d\tau_0(\mathfrak{l}_i \times \mathfrak{l}_i) \subset \mathfrak{a}_i$$

and

$$d\tau(L_1, L_2) = L_1 T_2(L_2) - L_2 T_1(L_1),$$

which in particular proves condition (i). It remains to check the last two conditions in (iii). By the above we have

$$\langle \alpha_0 \wedge \tau \rangle_{\mathfrak{a}}(L_1, L'_1, L_2) = \langle \alpha_0(L_1, L'_1), T_2(L_2) \rangle_{\mathfrak{a}} \quad (22)$$

and

$$\begin{aligned} \frac{1}{2} \langle d\tau \wedge \tau \rangle_{\mathfrak{a}}(L_1, L'_1, L_2) &= \frac{1}{2} \left(\langle d\tau(L_1, L'_1), T_2(L_2) \rangle_{\mathfrak{a}} \right. \\ &\quad - \langle L_1 T_2(L_2), \tau(L'_1) \rangle_{\mathfrak{a}} + \langle L_2 T_1(L_1), \tau(L'_1) \rangle_{\mathfrak{a}} \\ &\quad \left. + \langle L'_1 T_2(L_2), \tau(L_1) \rangle_{\mathfrak{a}} - \langle L_2 T_1(L'_1), \tau(L_1) \rangle_{\mathfrak{a}} \right) \\ &= \frac{1}{2} \left(\langle d\tau(L_1, L'_1), T_2(L_2) \rangle_{\mathfrak{a}} + \langle T_2(L_2), L_1 \tau(L'_1) - L'_1 \tau(L_1) \rangle_{\mathfrak{a}} \right) \\ &\quad - \langle L_2 \tau(L'_1), T_1(L_1) \rangle_{\mathfrak{a}} \\ &= \langle d\tau(L_1, L'_1), T_2(L_2) \rangle_{\mathfrak{a}} + \langle d\tau(L'_1, L_2), T_1(L_1) \rangle_{\mathfrak{a}}. \end{aligned} \quad (23)$$

It follows that

$$\begin{aligned} \gamma(L_1, L'_1, L_2) &= \langle (\alpha_0 + d\tau)(L_1, L'_1), T_2(L_2) \rangle_{\mathfrak{a}} + \langle d\tau(L'_1, L_2), T_1(L_1) \rangle_{\mathfrak{a}} \\ &= \langle \alpha(L_1, L'_1), T_2(L_2) \rangle_{\mathfrak{a}} + \langle \alpha(L'_1, L_2), T_1(L_1) \rangle_{\mathfrak{a}}. \end{aligned}$$

Changing the roles of the indices we also obtain

$$\gamma(L_2, L'_2, L_1) = \langle \alpha(L_2, L'_2), T_1(L_1) \rangle_{\mathfrak{a}} + \langle \alpha(L'_2, L_1), T_2(L_2) \rangle_{\mathfrak{a}}.$$

This finishes the proof of (iii).

Vice versa, let us assume that $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ satisfies the conditions (i)-(iii) for certain maps T_1, T_2 . We set

$$\tau := T_1 + T_2, \quad \alpha_0 := \alpha - d\tau, \quad \text{and} \quad \gamma_0 = \gamma - \langle (\alpha_0 + \frac{1}{2}d\tau) \wedge \tau \rangle_{\mathfrak{a}}.$$

Then, by Proposition 3.1 the algebras $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ and $\mathfrak{d}_{\alpha_0, \gamma_0}(\mathfrak{a}, \mathfrak{l}, \rho)$ are extension equivalent. By the remark at the beginning of the proof it is sufficient to show that α_0 and γ_0 are diagonal. That α_0 is diagonal is obvious. This allows us to repeat the computations (22) and (23) in order to obtain

$$\langle (\alpha_0 + \frac{1}{2}d\tau) \wedge \tau \rangle_{\mathfrak{a}}(L_1, L'_1, L'_2) = \langle \alpha(L_1, L'_1), T_2(L'_2) \rangle_{\mathfrak{a}} + \langle \alpha(L'_1, L'_2), T_1(L_1) \rangle_{\mathfrak{a}}.$$

By assumption, the right hand side is equal to $\gamma(L_1, L'_1, L'_2)$. It follows that

$$\gamma_0(L_1, L'_1, L'_2) = 0.$$

Changing the roles of the indices, we eventually see that γ_0 is diagonal. This finishes the proof of the proposition. $\square$

We conclude this section with a description of the isomorphism classes of non-abelian indecomposable metric Lie algebras with the property that $\mathfrak{g}'/\mathfrak{z}(\mathfrak{g})$ is abelian. Let us fix $\mathfrak{l}$, $\mathfrak{a}$, and the inner product $\langle \cdot, \cdot \rangle_{\mathfrak{a}}$. We consider the set $\text{Hom}(\mathfrak{l}, \mathfrak{so}(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}}))$ of all orthogonal representations of $\mathfrak{l}$ on $\mathfrak{a}$. If $\rho \in \text{Hom}(\mathfrak{l}, \mathfrak{so}(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}}))$ is fixed we denote the corresponding $\mathfrak{l}$ -module by $\mathfrak{a}_{\rho}$. The group $G := GL(\mathfrak{l}) \times O(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ acts on $\text{Hom}(\mathfrak{l}, \mathfrak{so}(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}}))$ by $(S, U)\rho := \text{Ad}(U) \circ \rho \circ S^{-1}$, on $C^q(\mathfrak{l}, \mathfrak{a})$ by $(S, U)\alpha := U(S^{-1})^*\alpha$, and on $\bigwedge^3 \mathfrak{l}^*$ by $(S, U)\gamma := (S^{-1})^*\gamma$. These actions are compatible with the differential d , the product $\langle \cdot \wedge \cdot \rangle_{\mathfrak{a}}$, and the action (16). In particular, if $g \in G$ and $\alpha \in Z_C^2(\mathfrak{l}, \mathfrak{a}_{\rho})$, then $g\alpha \in Z_C^2(\mathfrak{l}, \mathfrak{a}_{g\rho})$. We obtain an induced action of G on the disjoint union

$$\coprod_{\rho \in \text{Hom}(\mathfrak{l}, \mathfrak{so}(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}}))} \mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}_{\rho}) .$$

Let $\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}_{\rho})_0 \subset \mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}_{\rho})$ be the subset corresponding to all extension equivalence classes of regular indecomposable $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$. The relevant conditions for (α, γ) are given in Lemma 2.1 and Proposition 4.1. Then the set

$$\coprod_{\rho \in \text{Hom}(\mathfrak{l}, \mathfrak{so}(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}}))} \mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}_{\rho})_0$$

is G -invariant. Combining Proposition 2.2 and Lemma 3.1 with Corollary 3.1 we obtain

Proposition 4.2 *Fix $\mathfrak{l}$, $\mathfrak{a}$, and an inner product $\langle \cdot, \cdot \rangle_{\mathfrak{a}}$. We consider the class $\mathcal{A}(\mathfrak{l}, \mathfrak{a})$ of non-abelian indecomposable metric Lie algebras $\mathfrak{g}$ satisfying*

1. $\mathfrak{g}'/\mathfrak{z}(\mathfrak{g})$ is abelian and isomorphic to $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ as a semi-Euclidean vector space,
2. $\dim \mathfrak{z}(\mathfrak{g}) = \dim \mathfrak{l}$.

Then the set of isomorphism classes of $\mathcal{A}(\mathfrak{l}, \mathfrak{a})$ is in bijective correspondence with the orbit space of the action of $G = GL(\mathfrak{l}) \times O(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ on

$$\coprod_{\rho \in \text{Hom}(\mathfrak{l}, \mathfrak{so}(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}}))} \mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}_{\rho})_0 .$$

5 The case of maximal isotropic centre

In Section 2 we proved that any indecomposable metric Lie algebra of signature (p, q) , $p \leq q$, with non-trivial maximal isotropic centre is isomorphic to a regular metric Lie algebra $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ for a Euclidean vector space $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$. Therefore we specialize now some of the results from the previous sections to the case where $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ is Euclidean.

Remark 5.1 If $\mathfrak{a}$ is Euclidean the nil-subspace $\mathfrak{a}^{(1)}$ coincides with the space $\mathfrak{a}^{\mathfrak{l}}$ of invariants of the representation ρ . By Remark 3.2 we have

$$\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}) \cong \mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}^{\mathfrak{l}}).$$

Each extension equivalence class has a representative $\mathfrak{d}_{\alpha,\gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ such that $\alpha \in Z_C^2(\mathfrak{l}, \mathfrak{a}^{\mathfrak{l}})$. Furthermore, we have

$$\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}^{\mathfrak{l}}) = \left(Z_C^2(\mathfrak{l}, \mathfrak{a}^{\mathfrak{l}}) \times \Lambda^3 \mathfrak{l}^* \right) / C^1(\mathfrak{l}, \mathfrak{a}^{\mathfrak{l}}) = \left(\Lambda_C^2(\mathfrak{l}^*, \mathfrak{a}^{\mathfrak{l}}) \times \Lambda^3 \mathfrak{l}^* \right) / C^1(\mathfrak{l}, \mathfrak{a}^{\mathfrak{l}})$$

where $\Lambda_C^2(\mathfrak{l}^*, \mathfrak{a}^{\mathfrak{l}})$ denotes the space of 2-forms on $\mathfrak{l}^*$ with values in $\mathfrak{a}^{\mathfrak{l}}$ satisfying Equation (2). Here the action of $C^1(\mathfrak{l}, \mathfrak{a}^{\mathfrak{l}})$ on $\Lambda_C^2(\mathfrak{l}^*, \mathfrak{a}^{\mathfrak{l}}) \times \Lambda^3 \mathfrak{l}^*$ is given by

$$(\alpha, \gamma)\tau = (\alpha, \gamma + \langle \alpha \wedge \tau \rangle_{\mathfrak{a}}).$$

In particular, the orbit of $(\alpha, \gamma) \in \Lambda_C^2(\mathfrak{l}^*, \mathfrak{a}^{\mathfrak{l}}) \times \Lambda^3 \mathfrak{l}^*$ is an affine subspace of $\Lambda_C^2(\mathfrak{l}^*, \mathfrak{a}^{\mathfrak{l}}) \times \Lambda^3 \mathfrak{l}^*$ with associated vector space $\langle \alpha \wedge C^1(\mathfrak{l}, \mathfrak{a}^{\mathfrak{l}}) \rangle_{\mathfrak{a}} \subset \Lambda^3 \mathfrak{l}^*$. Hence

$$\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}^{\mathfrak{l}}) \cong \coprod_{\alpha \in \Lambda_C^2(\mathfrak{l}^*, \mathfrak{a}^{\mathfrak{l}})} \Lambda^3 \mathfrak{l}^* / \langle \alpha \wedge C^1(\mathfrak{l}, \mathfrak{a}^{\mathfrak{l}}) \rangle_{\mathfrak{a}}. \quad (24)$$

Since $\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}^{\mathfrak{l}})$ only depends on $\dim \mathfrak{l}$ and $\dim \mathfrak{a}^{\mathfrak{l}}$ we will denote it by $\mathcal{H}_C^2(\dim \mathfrak{l}, \dim \mathfrak{a}^{\mathfrak{l}})$.

Proposition 5.1 *Assume that $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ is Euclidean and $\alpha \in \Lambda_C^2(\mathfrak{l}^*, \mathfrak{a}^{\mathfrak{l}})$. We also assume that $\mathfrak{d}_{\alpha,\gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is not two-dimensional (the only two-dimensional algebra of this form is abelian and decomposable).*

Then $\mathfrak{d}_{\alpha,\gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is decomposable if and only if

(i) there are decompositions

$$\mathfrak{a} = \mathfrak{a}_1 \oplus \mathfrak{a}_2, \quad \mathfrak{l} = \mathfrak{l}_1 \oplus \mathfrak{l}_2$$

satisfying $\mathfrak{a}_1 \perp \mathfrak{a}_2$ and $\mathfrak{l}_i \oplus \mathfrak{a}_i \neq 0$, $i = 1, 2$, and representations ρ_i of $\mathfrak{l}_i$ on $\mathfrak{a}_i$ such that

$$\rho = \rho_1 \oplus \rho_2,$$

(ii) the cocycle $\alpha \in \Lambda_C^2(\mathfrak{l}^, \mathfrak{a}^{\mathfrak{l}})$ satisfies*

$$\alpha(\mathfrak{l}_1 \times \mathfrak{l}_1) \subset \mathfrak{a}_1, \quad \alpha(\mathfrak{l}_2 \times \mathfrak{l}_2) \subset \mathfrak{a}_2, \quad \alpha(\mathfrak{l}_1 \times \mathfrak{l}_2) = 0,$$

(iii) there are linear maps $T_1 : \mathfrak{l}_1 \rightarrow \mathfrak{a}_2$ and $T_2 : \mathfrak{l}_2 \rightarrow \mathfrak{a}_1$ such that

$$\gamma(L_1, L'_1, L_2) = \langle \alpha(L_1, L'_1), T_2 L_2 \rangle_{\mathfrak{a}}, \quad \gamma(L_2, L'_2, L_1) = \langle \alpha(L_2, L'_2), T_1 L_1 \rangle_{\mathfrak{a}}$$

for all $L_1, L'_1 \in \mathfrak{l}_1$ and $L_2, L'_2 \in \mathfrak{l}_2$.

Proof. If Conditions (i), (ii), (iii) are satisfied, then $\mathfrak{d}_{\alpha,\gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is decomposable by Proposition 4.1.

Assume now that $\mathfrak{d}_{\alpha,\gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is decomposable. If $\mathfrak{d}_{\alpha,\gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is regular, then we can again apply Proposition 4.1 and the assertion follows since

$$\alpha(L_1, L_2) = L_1 T_2 L_2 - L_2 T_1 L_1 \in \mathfrak{a}^{\mathfrak{l}}$$

implies $\alpha(L_1, L_2) = 0$ for $L_1 \in \mathfrak{l}_1$ and $L_2 \in \mathfrak{l}_2$. If $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is not regular then there exist elements $A_0 \in \mathfrak{a}$, $L_0 \in \mathfrak{l}$, $A_0 + L_0 \neq 0$ which satisfy Equations (3). If $L_0 = 0$, then

$$\mathfrak{a}_1 = \mathbb{R}A_0, \quad \mathfrak{a}_2 = (\mathbb{R}A_0)^\perp, \quad \mathfrak{l}_1 = 0, \quad \mathfrak{l}_2 = \mathfrak{l}, \quad T_1 = T_2 = 0$$

satisfy Conditions (i), (ii), (iii). For $L_0 \neq 0$ we can take

$$\mathfrak{a}_1 = 0, \quad \mathfrak{a}_2 = \mathfrak{a}, \quad \mathfrak{l}_1 = \mathbb{R}L_0, \quad T_1(L_0) = A_0, \quad T_2 = 0$$

and choose for $\mathfrak{l}_2$ any complement of $\mathbb{R}L_0$ in $\mathfrak{l}$. This decomposition of $\mathfrak{l} \oplus \mathfrak{a}$ is non-trivial in the sense of Condition (i) since by assumption $\mathfrak{d}_{\alpha, \gamma}(\mathfrak{a}, \mathfrak{l}, \rho)$ is at least three-dimensional. $\square$

Remark 5.2 Let $\mathfrak{l}$ be an abelian Lie algebra of dimension l and $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ a Euclidean space of dimension n . Recall that $GL(\mathfrak{l}) \times O(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ acts on $\text{Hom}(\mathfrak{l}, \mathfrak{so}(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}}))$ by

$$(S, U)\rho(\cdot) = U \circ \rho(S^{-1}(\cdot)) \circ U^{-1}.$$

The orbit space of this action equals

$$\begin{aligned} \text{Hom}(\mathfrak{l}, \mathfrak{so}(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})) / GL(\mathfrak{l}) \times O(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}}) &= (\mathfrak{l}^*)^m / (GL(\mathfrak{l}) \times \mathfrak{S}_m \ltimes (\mathbb{Z}_2)^m) \\ &= \coprod_{k \leq l} \mathbb{G}_{k, m} / (\mathfrak{S}_m \ltimes (\mathbb{Z}_2)^m), \end{aligned}$$

where $m = \lfloor \frac{n}{2} \rfloor$, $\mathbb{G}_{k, m}$ denotes the Grassmannian and $\mathfrak{S}_m$ the symmetric group. Here we first identify the orbit of $\rho \in \text{Hom}(\mathfrak{l}, \mathfrak{so}(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}}))$ with the $GL(\mathfrak{l}) \times \mathfrak{S}_m \ltimes (\mathbb{Z}_2)^m$ -orbit of the weights $\lambda = (\lambda^1, \dots, \lambda^m) \in (\mathfrak{l}^*)^m$ of ρ and then with the $\mathfrak{S}_m \ltimes (\mathbb{Z}_2)^m$ -orbit of $\text{span}\{\lambda(L) \mid L \in \mathfrak{l}\}$.

Example 5.1 Consider the abelian Lie algebra $\mathfrak{l} = \mathbb{R}^l$ and let $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ be the Euclidean space $\mathbb{R}^{2m} \oplus \mathbb{R}^k$. Choose $\lambda = (\lambda^1, \dots, \lambda^m) \in (\mathfrak{l}^*)^m$. Define an orthogonal representation ρ of $\mathfrak{l}$ on $\mathfrak{a}$ by

$$\rho(L) = \begin{pmatrix} 0 & -\text{diag}(\lambda(L)) & 0 \\ \text{diag}(\lambda(L)) & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Furthermore, assume $\alpha \in \wedge_C^2(\mathbb{R}^l, \mathbb{R}^k)$, $\gamma \in \wedge^3 \mathbb{R}^l$ and define

$$\mathfrak{d}_{\alpha, \gamma}(m, k, l, \lambda) := \mathfrak{d}_{\alpha, \gamma}(\mathbb{R}^{2m+k}, \mathbb{R}^l, \rho).$$

Next we want to give a method which yields (in principle) a description of the isomorphism classes of indecomposable metric Lie algebras of signature $(l, n + l)$, $l > 0$, with maximal isotropic centre. By Theorem 2.1 this set is equal to the set of isomorphism classes in $\mathcal{A}(\mathfrak{l}, \mathfrak{a})$ for a fixed n -dimensional Euclidean space $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ and a fixed l -dimensional abelian Lie algebra $\mathfrak{l}$. By Proposition 4.2 the isomorphism classes in $\mathcal{A}(\mathfrak{l}, \mathfrak{a})$ are in bijective correspondence with the $GL(\mathfrak{l}) \times O(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$ -orbits of

$$\coprod_{\rho \in \text{Hom}(\mathfrak{l}, \mathfrak{so}(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}}))} \mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}_\rho)_0.$$

Note that in the case of Euclidean $\mathfrak{a}$ the set $\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}_\rho)_0$ consists of all elements in $\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}_\rho)$ which correspond to indecomposable metric Lie algebras. According to Remark 5.1 and Remark 5.2 this orbit space equals

$$\begin{aligned} & \coprod_{k=0}^n \left(\left(\coprod_{\substack{\rho \in \text{Hom}(\mathfrak{l}, \mathfrak{so}(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})) \\ \dim \mathfrak{a}_\rho^{\mathfrak{l}} = k}} \mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}_\rho)_0 \right) / O(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}}) \right) / GL(\mathfrak{l}) \\ &= \coprod_{\substack{(k, m) \\ k + 2m = n}} \left(\left((\mathfrak{l}^* \setminus 0)^m / \mathfrak{S}_m \times (\mathbb{Z}_2)^m \right) \times \left(\mathcal{H}_C^2(\mathfrak{l}, k) / O(k) \right) \right)_0 / GL(\mathfrak{l}), \end{aligned}$$

where the subscript 0 again denotes the subset of elements which correspond to indecomposable metric Lie algebras. By Proposition 5.1 indecomposability implies that $\alpha(\mathfrak{l}, \mathfrak{l}) = \mathfrak{a}_\rho^{\mathfrak{l}}$. In fact, this property is already a consequence of regularity. In particular, $\mathcal{H}_C^2(\mathfrak{l}, \mathfrak{a}_\rho)_0 = 0$ for $k = \dim \mathfrak{a}_\rho^{\mathfrak{l}} > l(l-1)/2$, $l = \dim \mathfrak{l}$.

Hence, if we want to determine the isomorphism classes in $\mathcal{A}(\mathfrak{l}, \mathfrak{a})$, then for all (k, m) with $k + 2m = n$ we have to

1. determine $\mathcal{H}_C^2(l, k)$ for $k \leq l(l-1)/2$ according to (24),
2. determine representatives $[\alpha_\iota, \gamma_\iota]$ (ι runs over an index set I) of those $(GL(\mathfrak{l}) \times O(k))$ -orbits in $\mathcal{H}_C^2(l, k)$ whose elements $[\alpha, \gamma]$ satisfy $\alpha(\mathfrak{l}, \mathfrak{l}) = \mathbb{R}^k$,
3. determine the stabilizer $G_\iota \subset GL(\mathfrak{l}) \times O(k)$ for each $[\alpha_\iota, \gamma_\iota]$,
4. determine the subset $\Lambda \subset (\mathfrak{l}^* \setminus 0)^m$ of those $\lambda \in (\mathfrak{l}^* \setminus 0)^m$ for which $\mathfrak{d}_{\alpha_\iota, \gamma_\iota}(m, k, l, \lambda)$ is indecomposable according to Proposition 5.1,
5. determine the set $\{\mathcal{O}_\theta \mid \theta \in \Theta_\iota\}$, of $\bar{G}_\iota \times \mathfrak{S}_m \times (\mathbb{Z}_2)^m$ -orbits in Λ , where $\bar{G}_\iota = \text{proj}_{GL(\mathfrak{l})} G_\iota \subset GL(\mathfrak{l})$.

Then each isomorphism class of indecomposable metric Lie algebras of signature $(l, n+l)$ with maximal isotropic centre is represented by one of the metric Lie algebras $\mathfrak{d}_{\alpha_\iota, \gamma_\iota}(m, k, l, \lambda)$, where $2m + k = n$. Here $\iota \in I$ and the $\bar{G}_\iota$ -orbit $\mathcal{O}_\theta$, $\theta \in \Theta_\iota$ of λ are uniquely determined.

Remark 5.3 The main obstruction (besides problems of bookkeeping and the lack of time) which prevents us from carrying out this procedure in general is the rather complicated geometry of 3-forms. Indeed, Step 2 involves the determination of the orbit spaces

$$\left(\bigwedge^3 \mathfrak{l}^* / \langle \alpha \wedge C^1(\mathfrak{l}, \mathbb{R}^k) \rangle \right) / \bar{G}_\alpha, \quad \alpha \in \bigwedge_C^2(\mathfrak{l}^*, \mathbb{R}^k) \text{ such that } \alpha(\mathfrak{l}, \mathfrak{l}) = \mathbb{R}^k,$$

where $\bar{G}_\alpha \subset GL(\mathfrak{l})$ denotes projection of the stabilizer of α in $GL(\mathfrak{l}) \times O(k)$ onto the first factor. This problem is already difficult in the case $\alpha = 0$ (which implies $k = 0$). To our knowledge, the relevant orbit space

$$\bigwedge^3 \mathfrak{l}^* / GL(\mathfrak{l})$$

is not known for $l \geq 10$. If $l \leq 8$, then it is finite; see [Gu 64], §35 for a classification over $\mathbb{C}$, compare also [H 00]. For dimension $l = 9$ we refer to [Kt 92] and the references cited therein. Here the orbit space is (generically) 3-dimensional.

Now we will apply this method for metrics with small index. Let us start with Lorentzian Lie algebras. We already know from Corollary 2.1 that any indecomposable non-simple Lorentzian Lie algebra has a maximal isotropic centre. Therefore our recipe reproduces the known classification of (at least 2-dimensional) indecomposable non-simple Lorentzian Lie algebras (see [M 85]). Any such Lie algebra is isomorphic to exactly one of the indecomposable Lorentzian algebras

$$\text{osc}(\lambda^1, \dots, \lambda^m) := \mathfrak{d}_{0,0}(m, 0, 1, \lambda), \quad 1 = \lambda^1 \leq \lambda^2 \leq \dots \leq \lambda^m,$$

where $(\lambda^1, \dots, \lambda^m) = \lambda(L)$ for a fixed basis L of $\mathfrak{l} = \mathbb{R}$.

Note, that the only simple Lorentzian Lie algebra is $\mathfrak{sl}(2, \mathbb{R})$ since the Killing forms of all other simple Lie algebras (up to the sign) do not have Lorentzian signature. This completes the classification of all indecomposable Lorentzian Lie algebras.

In the following we will fix a basis $L_1, \dots, L_l$ of $\mathfrak{l} = \mathbb{R}^l$, the dual basis $Z_1, \dots, Z_l$ in $\mathfrak{l}^*$, and an orthonormal basis $A_1, \dots, A_k$ of $\mathbb{R}^k$. For $\lambda \in (\mathfrak{l}^*)^m$ we define $(\lambda_j^i)_{j=1, \dots, l}^{i=1, \dots, m} = (\lambda^i(L_j))_{j=1, \dots, l}^{i=1, \dots, m}$ and denote by $\lambda^i = (\lambda_1^i, \dots, \lambda_l^i)$, $i = 1, \dots, m$, and $\lambda_j = (\lambda_j^1, \dots, \lambda_j^m)^\top$, $j = 1, \dots, l$ the rows and the columns of $(\lambda_j^i)_{j=1, \dots, l}^{i=1, \dots, m}$, respectively.

Theorem 5.1 *For $l = 2, 3$ each isomorphism class of indecomposable metric Lie algebras of signature $(l, l + n)$ with maximal isotropic centre is represented by a metric Lie algebra $\mathfrak{d}_{\alpha_l, \gamma_l}(m, k, l, \lambda)$, $\lambda \in \Lambda_l$, for exactly one combination of m, k, α_l, γ_l in the table below and for a suitable λ in the set $\Lambda_l \subset (\mathfrak{l}^* \setminus 0)^m$, which is also given in the table. In particular, we have $n = 2m + k$.*

The metric Lie algebras $\mathfrak{d}_{\alpha_l, \gamma_l}(m, k, l, \lambda)$ and $\mathfrak{d}_{\alpha_l, \gamma_l}(m, k, l, \bar{\lambda})$ are isomorphic if and only if

$$\vartheta_l(\lambda) = \vartheta_l(\bar{\lambda}) \pmod{\mathfrak{S}_m \times (\mathbb{Z}_2)^m}$$

for the map ϑ_l which is associated with the combination m, k, α_l, γ_l and listed in the table.

Conversely, all the metric Lie algebras $\mathfrak{d}_{\alpha_l, \gamma_l}(m, k, l, \lambda)$ with $l, m, k, \alpha_l, \gamma_l, \lambda \in \Lambda_l$ as in the table are indecomposable and of signature $(l, l + 2m + k)$ with maximal isotropic centre.

$$l = 2$$

$k = 0$ $m \geq 3$	$\alpha = 0, \gamma = 0$ $\Lambda = \left\{ \lambda \in (\mathfrak{l}^* \setminus 0)^m \mid \begin{array}{l} \{\lambda^i \mid i = 1, \dots, m\} \subset \mathfrak{l}^* \text{ is not contained in the} \\ \text{union of two 1-dimensional subspaces of } \mathfrak{l}^* \end{array} \right\}$ $\vartheta(\lambda) = \text{span}\{\lambda_1, \lambda_2\} \in \mathbb{G}_{2,m}$
-----------------------	--

$k = 1$ $m \geq 0$	$\alpha = Z_1 \wedge Z_2 \otimes A_1$, $\gamma = 0$ $\Lambda = (\mathfrak{l}^* \setminus 0)^m$ $\vartheta(\lambda) = (E_\lambda, \omega_\lambda)$, where $E_\lambda = \text{span}\{\lambda_1, \lambda_2\} \subset \mathbb{R}^m$, $\omega_\lambda = [\lambda_1 \wedge \lambda_2] \in \wedge^2(E_\lambda)/\{\pm 1\}$
-----------------------	--

$$l = 3$$

$k = 0$ $m \geq 4$	$\alpha_1 = 0, \gamma_1 = 0$ $\Lambda_1 = \left\{ \lambda \in (\mathfrak{l}^* \setminus 0)^m \left \begin{array}{l} \{\lambda^i \mid i = 1, \dots, m\} \subset \mathfrak{l}^* \text{ is not contained} \\ \text{in the union of a 2-dimensional and a} \\ \text{1-dimensional subspace of } \mathfrak{l}^* \end{array} \right. \right\}$ $\vartheta_1(\lambda) = \text{span}\{\lambda_1, \lambda_2, \lambda_3\} \in \mathbb{G}_{3,m}$
$k = 0$ $m \geq 0$	$\alpha_2 = 0, \gamma_2 = Z_1 \wedge Z_2 \wedge Z_3$ $\Lambda_2 = (\mathfrak{l}^* \setminus 0)^m$ $\vartheta_2(\lambda) = (E_\lambda, \omega_\lambda)$, where $E_\lambda = \text{span}\{\lambda_1, \lambda_2, \lambda_3\} \subset \mathbb{R}^m$, $\omega_\lambda = \lambda_1 \wedge \lambda_2 \wedge \lambda_3 \in \wedge^3(E_\lambda)$
$k = 1$ $m \geq 2$	$\alpha = Z_1 \wedge Z_2 \otimes A_1$, $\gamma = 0$ $\Lambda = \left\{ \lambda \in (\mathfrak{l}^* \setminus 0)^m \mid \dim \text{span}\{\lambda^j \mid \lambda_3^j \neq 0\} > 1 \right\}$ $\vartheta(\lambda) = (E_\lambda, F_\lambda, \omega_\lambda)$, where $E_\lambda = \text{span}\{\lambda_1, \lambda_2, \lambda_3\} \subset \mathbb{R}^m$, $F_\lambda = \mathbb{R}\lambda_3 \subset E_\lambda$, $\omega_\lambda = [\lambda_1 \wedge \lambda_2 \bmod \mathbb{R}\lambda_3] \in \wedge^2(E_\lambda/F_\lambda)/\{\pm 1\}$
$k = 2$ $m \geq 0$	$\alpha = Z_1 \wedge Z_2 \otimes A_1 + Z_1 \wedge Z_3 \otimes A_2, \gamma = 0$ $\Lambda = (\mathfrak{l}^* \setminus 0)^m$ $\vartheta(\lambda) = (B_\lambda, v_\lambda) \bmod \mathbb{R}^*$, where $B_\lambda = \left((\lambda_2^i, \lambda_3^i) \cdot (\lambda_2^j, \lambda_3^j)^\top \right)_{j=1, \dots, m}^{i=1, \dots, m} \in \text{End}(\mathbb{R}^m)$, $v_\lambda = [\lambda_1] \in \text{coker } B_\lambda$, $c \in \mathbb{R}^*$ acts by $(B, v)c = (c^2 B, \frac{1}{c} v)$
$k = 3$ $m \geq 0$	$\alpha = Z_1 \wedge Z_2 \otimes A_1 + Z_1 \wedge Z_3 \otimes A_2 + Z_2 \wedge Z_3 \otimes A_3$, $\gamma = 0$ $\Lambda = (\mathfrak{l}^* \setminus 0)^m$ $\vartheta(\lambda) = \left(\lambda^i \cdot (\lambda^j)^\top \right)_{j=1, \dots, m}^{i=1, \dots, m} \in \text{End}(\mathbb{R}^m)$

Proof. We restrict ourselves to the case $l = 3$. We follow the recipe described above. First we note that for $l = 3$ Equation (2) is trivially satisfied. Then it is not hard to see that each class in $\mathcal{H}_G^2(3, k)$ is represented by exactly one element

$$(\alpha, \gamma) \in \left(\{0\} \times \Lambda^3 \mathfrak{l}^* \right) \cup \left((\Lambda^2(\mathfrak{l}^*, \mathbb{R}^k) \setminus \{0\}) \times \{0\} \right) \subset \Lambda^2(\mathfrak{l}^*, \mathbb{R}^k) \oplus \Lambda^3 \mathfrak{l}^*.$$

The group $GL(\mathfrak{l}) \times O(k)$ acts on this set of representatives. We have to determine the orbits of this action and to choose representatives, where we may restrict ourselves to orbits of those (α, γ) which satisfy $\dim(\alpha(\mathfrak{l}, \mathfrak{l})) = k$. Obviously, for $k = 0$, i.e. $\alpha = 0$ we have two such orbits, one orbit consists of the element $(\alpha, \gamma) = (0, 0)$, the other one is characterized by $\gamma \neq 0$. For $k \geq 1$ we have $\alpha \neq 0$ and may therefore assume that $\gamma = 0$ holds. We claim that in this case there is only one orbit. The representation $\Lambda^2 \mathfrak{l}$ of $GL(\mathfrak{l})$ is equivalent to the representation ρ of $GL(\mathfrak{l})$ on $\mathfrak{l}^*$ given by $GL(\mathfrak{l}) \ni S \mapsto \det S \cdot (S^{-1})^* \in GL(\mathfrak{l}^*)$. In particular, $GL(\mathfrak{l})$ acts transitively on oriented bases in $\Lambda^2 \mathfrak{l}$. Therefore, given a basis $\sigma^1, \sigma^2, \sigma^3$ in $\Lambda^2 \mathfrak{l}$, a basis $A_1, \dots, A_k$ of $\mathbb{R}^k$ and a 2-form $\alpha \in \Lambda^2(\mathfrak{l}^*, \mathbb{R}^k)$ we find a map $S \in GL(\mathfrak{l})$ such that $(S^* \alpha)(\sigma^j) = \pm A_j$ for $1 \leq j \leq k$ and $(S^* \alpha)(\sigma^j) = 0$ for $k < j \leq 3$. Now it is obvious that there exists a map $U \in O(k)$ such that $(US^* \alpha)(\sigma^j) = A_j$ for $1 \leq j \leq k$ and $(US^* \alpha)(\sigma^j) = 0$ for $k < j \leq 3$, which proves the claim. In particular, we may choose (α, γ) as in the table. Since the proofs of the assertions on indecomposability and isomorphy are similar for different k we will give here only the one for $k = 2$, which is in some sense the most complicated one. First we determine the projection $\bar{G} = \text{proj}_{GL(\mathfrak{l})} G$ of the stabilizer $G \subset GL(\mathfrak{l}) \times O(2)$ of $(\alpha, \gamma) = (Z_1 \wedge Z_2 \otimes A_1 + Z_1 \wedge Z_3 \otimes A_2, 0)$ on $GL(\mathfrak{l})$:

$$\begin{aligned} \bar{G} &= \left\{ S \in GL(\mathfrak{l}) \left| \begin{array}{l} S^* \alpha(L_1, L_2), S^* \alpha(L_1, L_3) \text{ is an orthonormal basis of } \mathbb{R}^k, \\ S^* \alpha(L_2, L_3) = 0 \end{array} \right. \right\} \\ &= \left\{ S \in GL(\mathfrak{l}) \left| \begin{array}{l} S(\text{span}\{L_2, L_3\}) = \text{span}\{L_2, L_3\} \\ S|_{\text{span}\{L_2, L_3\}} \in c \cdot O(2) \text{ with respect to} \\ \text{the basis } \{L_2, L_3\}, \text{ where } c = |\det S| \end{array} \right. \right\}. \end{aligned} \quad (25)$$

Next we determine the set $\Lambda \subset (\mathfrak{l}^* \setminus 0)^m$ of those $\lambda \in (\mathfrak{l}^* \setminus 0)^m$ for which $\mathfrak{d}_{\alpha, 0}(m, 2, 3, \lambda)$ is indecomposable. Assume that $\mathfrak{d}_{\alpha, 0}(m, 2, 3, \lambda)$ is decomposable. By Proposition 5.1 we have decompositions $\mathfrak{l} = \mathbb{R}^3 = \mathfrak{l}_1 \oplus \mathfrak{l}_2$ and $\mathfrak{a} = \mathbb{R}^{2m+k} = \mathfrak{a}_1 \oplus \mathfrak{a}_2$, at least one of them non-trivial, such that $\rho = \rho_1 \oplus \rho_2$ holds for the representation ρ defined by $\lambda \in (\mathfrak{l}^* \setminus 0)^m$ and α satisfies $\alpha(\mathfrak{l}_1 \times \mathfrak{l}_1) \subset \mathfrak{a}_1$, $\alpha(\mathfrak{l}_2 \times \mathfrak{l}_2) \subset \mathfrak{a}_2$, $\alpha(\mathfrak{l}_1 \times \mathfrak{l}_2) = 0$. Since $\ker \alpha \subset \Lambda^2(\mathfrak{l}^*, \mathfrak{a})$ is only 1-dimensional the condition $\alpha(\mathfrak{l}_1 \times \mathfrak{l}_2) = 0$ implies that the decomposition of $\mathfrak{l}$ is trivial. Hence, we may assume $\mathfrak{l}_1 = \mathfrak{l}$ and $\mathfrak{l}_2 = 0$. Thus the decomposition of $\mathfrak{a}$ is non-trivial, i.e. $\mathfrak{a}_2 \neq 0$. Now $\alpha(\mathfrak{l}_1 \times \mathfrak{l}_1) \subset \mathfrak{a}_1$ implies $\alpha(\mathfrak{l} \times \mathfrak{l}) = \mathbb{R}^k \subset \mathfrak{a}_1$. On the other hand $\mathfrak{l}_2 = 0$ yields $0 \neq \mathfrak{a}_2 \subset \mathfrak{a}^\mathfrak{l}$ which is a contradiction to $\lambda^j \neq 0$ for $j = 1, \dots, m$. Consequently, $\Lambda = (\mathfrak{l}^* \setminus 0)^m$.

Finally, we determine the $\bar{G}$ -orbits in $(\mathfrak{l}^* \setminus 0)^m$. From (25) we know that $(\lambda_1, \lambda_2, \lambda_3)$ and $(\bar{\lambda}_1, \bar{\lambda}_2, \bar{\lambda}_3)$ are in the same orbit if and only if there is a map $S' \in O(2)$ and a real number $c \in \mathbb{R}^*$ such that $c(\lambda_2, \lambda_3) \cdot S' = (\bar{\lambda}_2, \bar{\lambda}_3)$ and $\lambda_1 = c\bar{\lambda}_1 \text{ mod } \text{span}\{\lambda_2, \lambda_3\}$. Now we use the following general fact, which is true for arbitrary homomorphisms $A, \bar{A} : \mathbb{R}^N \rightarrow \mathbb{R}^m$:

$$\exists s \in O(N) : \bar{A} = A \cdot s \quad \Leftrightarrow \quad A A^* = \bar{A} \bar{A}^* : \mathbb{R}^m \longrightarrow \mathbb{R}^m.$$

We apply this for $A = (\lambda_j^i)_{j=2,3}^{i=1,\dots,m}$ and $\bar{A} = (\bar{\lambda}_j^i)_{j=2,3}^{i=1,\dots,m}$ and use the notation $B_\lambda = (\lambda_j^i)_{j=2,3}^{i=1,\dots,m} ((\lambda_j^i)_{j=2,3}^{i=1,\dots,m})^\top$. Note that $\text{span}\{\lambda_2, \lambda_3\} = \text{im}(\lambda_j^i)_{j=2,3}^{i=1,\dots,m} = \text{im } B_\lambda$. Now we can say that $(\lambda_1, \lambda_2, \lambda_3)$ and $(\bar{\lambda}_1, \bar{\lambda}_2, \bar{\lambda}_3)$ are in the same $\bar{G}$ -orbit if and only if there exists a real number $c \in \mathbb{R}^*$ such that $c^2 B_\lambda = B_{\bar{\lambda}}$ and $[\lambda_1] = c[\bar{\lambda}_1] \in \text{coker } B_\lambda (= \text{coker } B_{\bar{\lambda}})$. $\square$

Remark 5.4 Based on the classification results mentioned in Remark 5.3, it should be not too difficult to obtain a classification of indecomposable metric Lie algebras with maximal isotropic centre of signature (l, l) , $2 \leq l \leq 9$. The isomorphism classes of these metric Lie algebras are parametrized by those $[\gamma] \in \bigwedge^3 \mathfrak{l}^*/GL(\mathfrak{l})$ which are not represented by elements of the form $\gamma = \gamma_1 + \gamma_2$, $\gamma_i \in \bigwedge^3 \mathfrak{l}_i^*$ for a nontrivial decomposition $\mathfrak{l} = \mathfrak{l}_1 \oplus \mathfrak{l}_2$, via

$$[\gamma] \longmapsto \mathfrak{d}_{0,\gamma}(0, 0, l, 0) .$$

In fact, it is easy to carry out the classification for $l \leq 5$. There are no such algebras for $l = 2, 4$ and exactly one for $l = 3, 5$. Moreover, it can be shown that there are exactly two for $l = 6$.

6 Metric Lie algebras of index 2

Example 6.1 For a map $A \in \mathfrak{so}(p, q)$ we consider $\mathfrak{d}_A(\mathbb{R}^{p,q}, \mathbb{R}) := \mathfrak{d}_{0,0}(\mathbb{R}^{p,q}, \mathbb{R}, \rho)$, where ρ is defined by $\rho(1) = A$. Recall that $\mathfrak{d}_A(\mathbb{R}^{p,q}, \mathbb{R})$ is a metric Lie algebra of signature $(p+1, q+1)$ which can be described as follows: $\mathfrak{d}_A(\mathbb{R}^{p,q}, \mathbb{R})$ is isomorphic to $\mathbb{R}^{1,1} \oplus \mathbb{R}^{p,q}$ as a pseudo-Euclidean vector space, there exists a basis L, Z of $\mathbb{R}^{1,1}$ with $\langle L, L \rangle = \langle Z, Z \rangle = 0$ and $\langle L, Z \rangle = 1$ such that $Z \in \mathfrak{z}(\mathfrak{d}_A(\mathbb{R}^{p,q}, \mathbb{R}))$ and

$$[X_1, X_2] = \langle AX_1, X_2 \rangle Z, \quad [L, X] = A(X)$$

for all $X \in \mathbb{R}^{p,q}$.

Example 6.2 Consider the metric Lie algebra $\mathfrak{osc}(\lambda_1, \lambda_2) = \mathfrak{d}_{0,0}(m, 0, 2, \lambda)$ of signature $(2, 2+2m)$ which was defined in the previous section, where $\lambda_1 = \lambda(L_1)$, $\lambda_2 = \lambda(L_2) \in \mathbb{R}^m$ for a fixed basis L_1, L_2 of $\mathbb{R}^2$. Recall that this algebra can be described in the following way: $\mathfrak{osc}(\lambda_1, \lambda_2)$ for $\lambda_1, \lambda_2 \in \mathbb{R}^m$ is isomorphic to $\mathbb{R}^{2,2} \oplus \mathbb{R}^{2m}$ as a pseudo-Euclidean vector space, there exists a basis L_1, L_2, Z_1, Z_2 of $\mathbb{R}^{2,2}$ with

$$\langle L_i, L_j \rangle = \langle Z_i, Z_j \rangle = 0, \quad \langle Z_i, L_j \rangle = \delta_{ij}$$

for $i, j = 1, 2$ and an orthonormal basis $X_1, \dots, X_m, Y_1, \dots, Y_m$ of $\mathbb{R}^{2m}$ such that

$$\begin{aligned} Z_1, Z_2 &\in \mathfrak{z}(\mathfrak{osc}(\lambda_1, \lambda_2)) \\ [X_j, Y_k] &= \delta_{jk}(\lambda_1^j Z_1 + \lambda_2^j Z_2), \quad [X_j, X_k] = 0, \quad [Y_j, Y_k] = 0 \\ [L_i, X_j] &= \lambda_i^j Y_j, \quad [L_i, Y_j] = -\lambda_i^j X_j \\ [L_1, L_2] &= 0 \end{aligned}$$

for $i = 1, 2$ and $j, k = 1, \dots, m$,

Example 6.3 Now consider the metric Lie algebras $\mathfrak{d}(\lambda_1, \lambda_2) := \mathfrak{d}_{\alpha,0}(m, 1, 2, \lambda)$ for a fixed $\alpha \neq 0$ which have signature $(2, 2 + 2m + 1)$ and were defined in the previous section, too. Here $\lambda_1 = \lambda(L_1)$, $\lambda_2 = \lambda(L_2) \in \mathbb{R}^m$ for a fixed basis L_1, L_2 of $\mathbb{R}^2$ which is chosen in such a way that $\alpha(L_1, L_2)$ is a unit vector. This algebra can be described as follows: $\mathfrak{d}(\lambda_1, \lambda_2)$ for $\lambda_1, \lambda_2 \in \mathbb{R}^m$ is isomorphic to $\mathbb{R}^{2,2} \oplus \mathbb{R}^{2m+1}$ as a pseudo-Euclidean vector space, there exists a basis L_1, L_2, Z_1, Z_2 of $\mathbb{R}^{2,2}$ with

$$\langle L_i, L_j \rangle = \langle Z_i, Z_j \rangle = 0, \quad \langle Z_i, L_j \rangle = \delta_{ij}$$

for $i, j = 1, 2$ and an orthonormal basis $X_1, \dots, X_m, Y_1, \dots, Y_m, A_0$ of $\mathbb{R}^{2m+1}$ such that

$$\begin{aligned} Z_1, Z_2 &\in \mathfrak{z}(\mathfrak{d}(\lambda_1, \lambda_2)) \\ [A_0, X_j] &= [A_0, Y_j] = 0 \\ [A_0, L_1] &= Z_2, \quad [A_0, L_2] = -Z_1 \\ [X_j, Y_k] &= \delta_{jk}(\lambda_1^j Z_1 + \lambda_2^j Z_2), \quad [X_j, X_k] = 0, \quad [Y_j, Y_k] = 0 \\ [L_i, X_j] &= \lambda_i^j Y_j, \quad [L_i, Y_j] = -\lambda_i^j X_j \\ [L_1, L_2] &= A_0 \end{aligned}$$

for $i = 1, 2$ and $j, k = 1, \dots, m$

Theorem 6.1 *Let $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ be an indecomposable metric Lie algebra of signature $(2, q)$. If $\mathfrak{g}$ is simple, then $\mathfrak{g}$ is isomorphic to $\mathfrak{sl}(2, \mathbb{R})$ and $\langle \cdot, \cdot \rangle$ is a multiple of the Killing form. If $\mathfrak{g}$ is not simple, then the centre $\mathfrak{z}(\mathfrak{g})$ of $\mathfrak{g}$ is one- or two-dimensional and we are in one of the following cases.*

1. *If $\dim \mathfrak{z}(\mathfrak{g}) = 1$, then q is even and $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is isomorphic to $\mathfrak{d}_{L_2}(\mathbb{R}^{1,1}, \mathbb{R})$ if $q = 2$, or to exactly one of the spaces $\mathfrak{d}_{L_2, \lambda}(\mathbb{R}^{1, q-1}, \mathbb{R})$ if $q \geq 4$, where*

$$L_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad L_{2, \lambda} = \begin{pmatrix} L_2 & 0 & 0 \\ 0 & 0 & -\text{diag}(\lambda) \\ 0 & \text{diag}(\lambda) & 0 \end{pmatrix},$$

$\lambda = (\lambda^1, \lambda^2, \dots, \lambda^r)$ with $0 < \lambda^1 \leq \dots \leq \lambda^r$, $r = q/2 - 1$.

2. *If $\dim \mathfrak{z}(\mathfrak{g}) = 2$ and $\dim \mathfrak{g}$ is even, then $q = 2m + 2$ with $m \geq 3$ and $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is isomorphic to $\mathfrak{osc}(\lambda_1, \lambda_2)$ for some $\lambda_1, \lambda_2 \in \mathbb{R}^m$. There is no index $j \in \{1, \dots, m\}$ such that $\lambda_1^j = \lambda_2^j = 0$ and the set $\{(\lambda_i^1, \lambda_i^2) \mid i = 1, \dots, m\}$ is not contained in the union of two 1-dimensional subspaces. Two such Lie algebras $\mathfrak{osc}(\lambda_1, \lambda_2)$ and $\mathfrak{osc}(\bar{\lambda}_1, \bar{\lambda}_2)$ are isomorphic if and only if*

$$\text{span}\{\lambda_1, \lambda_2\} = \text{span}\{\bar{\lambda}_1, \bar{\lambda}_2\} \quad \text{mod } \mathfrak{S}_m \times (\mathbb{Z}_2)^m.$$

3. *If $\dim \mathfrak{z}(\mathfrak{g}) = 2$ and $\dim \mathfrak{g}$ is odd, then $q = 2m + 3$ for $m \geq 0$ and $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is isomorphic to $\mathfrak{d}(\lambda_1, \lambda_2)$ for some $\lambda_1, \lambda_2 \in \mathbb{R}^m$. There is no index $j \in \{1, \dots, m\}$ such that $\lambda_1^j = \lambda_2^j = 0$. Two such Lie algebras $\mathfrak{d}(\lambda_1, \lambda_2)$ and $\mathfrak{d}(\bar{\lambda}^1, \bar{\lambda}^2)$ are isomorphic if and only if*

$$(\text{span}\{\lambda_1, \lambda_2\}, \pm \lambda_1 \wedge \lambda_2) = (\text{span}\{\bar{\lambda}_1, \bar{\lambda}_2\}, \pm \bar{\lambda}_1 \wedge \bar{\lambda}_2) \quad \text{mod } \mathfrak{S}_m \times (\mathbb{Z}_2)^m.$$

Conversely, all metric Lie algebras $\mathfrak{d}_{L_2}(\mathbb{R}^{1,1}, \mathbb{R})$, $\mathfrak{d}_{L_{2,\lambda}}(\mathbb{R}^{1,q-1}, \mathbb{R})$, $\mathfrak{osc}(\lambda_1, \lambda_2)$, $\mathfrak{d}(\lambda_1, \lambda_2)$ are indecomposable and of signature $(2, q)$ if λ , λ_1 , λ_2 satisfy the conditions described above.

Proof. If $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is simple, then it is isomorphic to $\mathfrak{sl}(2, \mathbb{R})$, because there is no other simple Lie algebra whose Killing form has signature $(2, q)$. If $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is not simple, then by Proposition 2.1 there exist an abelian Lie algebra $\mathfrak{l}$, $\dim \mathfrak{l} = \dim \mathfrak{z}(\mathfrak{g}) \in \{1, 2\}$, a semi-Euclidean vector space $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}})$, an orthogonal representation ρ of $\mathfrak{l}$ on $\mathfrak{a}$, and a cocycle $\alpha \in Z^2(\mathfrak{l}, \mathfrak{a})$ satisfying (2), such that $\mathfrak{d}_{\alpha,0}(\mathfrak{a}, \mathfrak{l}, \rho)$ is regular and $(\mathfrak{g}, \langle \cdot, \cdot \rangle)$ is isomorphic to $\mathfrak{d}_{\alpha,0}(\mathfrak{a}, \mathfrak{l}, \rho)$.

Let us first consider the case $\dim \mathfrak{z}(\mathfrak{g}) = 1$. Then $\mathfrak{l} = \mathbb{R}$, $(\mathfrak{a}, \langle \cdot, \cdot \rangle_{\mathfrak{a}}) = \mathbb{R}^{1,q-1}$ and $\alpha = 0$. Hence, $(\mathfrak{g}, \langle \cdot, \cdot \rangle) = \mathfrak{d}_A(\mathbb{R}^{1,q-1}, \mathbb{R})$ as in Example 6.1. Since $\mathfrak{d}_A(\mathbb{R}^{1,q-1}, \mathbb{R})$ is regular we have $\ker A = 0$ by Lemma 2.1. By Corollary 3.2 regular metric Lie algebras $\mathfrak{d}_{A_1}(\mathbb{R}^{1,q-1}, \mathbb{R})$ and $\mathfrak{d}_{A_2}(\mathbb{R}^{1,q-1}, \mathbb{R})$ are isomorphic if and only if

$$\exists \mu \in \mathbb{R}^* \exists U \in O(1, q-1) : U A_1 U^{-1} = \mu A_2. \quad (26)$$

It is well-known that each equivalence class of bijective maps in $\mathfrak{so}(1, q-1)$ with respect to relation (26) is represented by exactly one of the maps L_2 (if $q = 2$) or $L_{2,\lambda}$ with $0 < \lambda^1 \leq \dots \leq \lambda^r$, $r = q/2 - 1$.

If $\dim \mathfrak{z}(\mathfrak{g}) = 2$, then the centre is maximal isotropic and the remaining assertions follow directly from Theorem 5.1. $\square$

References

- [BK 02] H. Baum, I. Kath, *Doubly Extended Lie Groups – Curvature, Holonomy and Parallel Spinors*. SFB 288 Preprint No. 543, 2002, see also Preprint arXiv:math.DG/0203189.
- [CP 70] M. Cahen, M. Parker, *Sur des classes d’espaces pseudo-riemanniens symétriques*. Bull. Soc. Math. Belg. **22** (1970), 339–354.
- [CP 80] M. Cahen, M. Parker, *Pseudo-Riemannian symmetric spaces*. Mem. Amer. Math. Soc. **24** (1980), no. 229.
- [CW 70] M. Cahen, N. Wallach, *Lorentzian symmetric spaces*. Bull. Amer. Math. Soc. **76** (1970), 585–591.
- [FS 96] J. M. Figueroa-O’Farrill, S. Stanciu, *On the structure of symmetric self-dual Lie algebras*. J. Math. Phys. **37** (1996), no. 8, 4121–4134.
- [Gr 98] A. N. Grishkov, *Orthogonal modules and nonlinear cohomologies*. Algebra and Logic **37** (1998), 294–306.
- [Gu 64] G. B. Gurevich, *Foundations of the Theory of Algebraic Invariants*. P. Noordhoff, Groningen, 1964.
- [H 00] N. Hitchin, *The geometry of three-forms in six and seven dimensions*. Preprint arXiv:math.DG/0010054, 2000.

- [Kt 92] A. A. Katanova, *Explicit form of certain multivector invariants*, in *Lie Groups, Their Discrete Subgroups and Invariant Theory*, pages 87–93. Advances in Soviet Mathematics, Vol. 8, AMS Providence, 1992.
- [M 85] A. Medina, *Groupes de Lie munis de métriques bi-invariantes*. Tohoku Math. J. (2) **37** (1985), no. 4, 405–421.
- [MR 85] A. Medina, Ph. Revoy, *Algèbres de Lie et produit scalaire invariant*. Ann. Sci. École Norm. Sup. (4) **18** (1985), no. 3, 553–561.
- [N 02] Th. Neukirchner, *Pseudo-Riemannian Symmetric Spaces*. Diplomarbeit, Humboldt-Universität zu Berlin, 2002.

Ines Kath, Martin Olbrich
 Mathematisches Institut der Georg-August-Universität Göttingen
 Bunsenstraße 3-5, D-37073 Göttingen, Germany
 email: kath@uni-math.gwdg.de, olbrich@uni-math.gwdg.de